

DIRECTIONS

SEVIN 50W is a wettable spray powder. Measure in level tablespoonfuls. Keep spray mixture well agitated. Do not mix with lime, Bordeaux Mixture or other alkaline materials. Do not mix with oil base sprays. Do not reuse container. Destroy when empty.

VEGETABLES

Mix 2 tablespoonfuls per gallon of water. Apply as a full wetting spray. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary unless shorter interval is specified below.

May be used up to day of harvest on the following: **BEANS:** Mexican bean beetle, leafhopper, cucumber beetle, Japanese beetle. **CORN:** corn rootworm adults, European corn borer, fall armyworm, corn earworm. Applications for corn earworms should be started when silks first appear. Repeat at 2 to 3-day intervals for corn earworms or until silks are dry. **CARROTS:** flea beetles, leafhoppers, harlequin bug. **CUCUMBER, SQUASH, PUMPKINS, MELONS:** pickleworm, melonworm, cucumber beetles, squash bug, flea beetle, leafhoppers. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

OKRA: corn earworm, stink bug. **POTATO:** Colorado potato beetle, flea beetle, leafhopper. Apply as a foliage application only. **TOMATO, EGGPLANT, PEPPER:** tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs and lygus bugs.

Do not apply within 3 days of harvest on the following: **BEETS, TURNIPS,** (do not apply within 14 days of harvest if tops are to be used as food or feed.) Leafhoppers. **BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI:** flea beetle, harlequin bug, armyworm, cabbage caterpillar, and corn earworm. **RADISHES:** flea beetle, leafhopper.

Do not apply within 14 days of harvest on the following: **LETTUCE (LEAF), SPINACH, SWISS CHARD:** armyworm, flea beetle, leafhopper.

LAWN INSECT CONTROL

Use one level cupful in 15 to 20 gallons of water for each 500 square feet of established lawn area for the control of: ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, **SOD WEBWORMS** (lawn moths) millipeds. Use **GARDEN HOSE SPRAYERS** or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately

after rain or watering. Repeat 2 to 3 weeks later if necessary.

ORNAMENTAL INSECT CONTROL

Ornamental plants, flowers, shrubs and shade trees including rose, carnation, gladiolus, zinnia, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not use on Boston ivy.

Use 2 level tablespoonfuls per gallon of spray (2 pounds per 100 gallons) when insects or their damage appear. Repeat weekly or as needed to control: apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf beetle, elm leaf aphid, flea beetle, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetles.

SMALL FRUIT INSECT CONTROL

Use sufficient spray to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary.

GRAPES

Use 2 tablespoonfuls per 1 gallon of water for European fruit lecanium, grape leaf folder, grape leafhopper, and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge from rolls, and as needed for leafhoppers. May be applied up to day of harvest.

STRAWBERRIES

For meadow spittlebug, leaf roller, strawberry weevil, use 2 tablespoonfuls per gallon of water. Make 3 to 5 post bloom summer sprays as needed. May be used up to 1 day before picking.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Packed by weight, not in volume. Product may settle in shipment.

USDA Reg. No. 3772-13

J-6309A-G

MANUFACTURED FOR

EARL MAY SEED & NURSERY CO.

SHENANDOAH, IOWA



SEVIN[★] WETTABLE POWDER

ACCEPTED
1-25-71
3772-13
UNION CARBIDE CORPORATION
PHOSPHORUS
POST OFFICE BOX 100
SPRINGFIELD, ILLINOIS
66101

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service

ACTIVE INGREDIENTS:	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	50%
INERT INGREDIENTS	50%
Total	100%

*SEVIN is a registered trade mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

CAUTION

KEEP OUT OF THE REACH OF CHILDREN.
HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Do Not Breathe Dust. Do Not Take Internally. Avoid Prolonged or Repeated Breathing of Dust. **Skin Contact May Be Harmful.** Avoid Skin Contact. Wash Hands Before Eating. **Avoid Contamination of Feed and Foodstuffs.** Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

NET WEIGHT 1 LB.

3772-13

01/25/71

1/1