

25 lbs.  
NET WEIGHT

# MAYTONE

## DANDELION

### WEED CONTROL PLUS FERTILIZER

#### GUARANTEED ANALYSIS

##### ACTIVE INGREDIENTS:

Diethylamine salt of 2,4-Dichlorophenoxyacetic acid*	0.72%
Dialkyl ether esters of silvex 2-(2,4,5-Trichlorophenoxy) propionic acid**	0.40%
Technical Calordane***	1.28%

##### NET WEIGHT INGREDIENTS

Relative to Herbicides	98.68%
Relative to Insecticide	99.72%

- \*Equivalent to 0.60% 2,4-Dichlorophenoxyacetic acid  
\*\*Equivalent to 0.40% 2-(2,4,5-Trichlorophenoxy) propionic acid  
\*\*\*Equivalent to 0.75% Caloridin, 7-Methoxy-2-methyl-2H-benzofuran-3-one compounds.

#### FERTILIZER

Total Nitrogen (N)	10%
7% Ammoniacal Nitrogen	
3% Nitrate Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	6%
From Superphosphate	
Soluble Potash (K <sub>2</sub> O)	4%
From Muriate of Potash	
Potential acidity 900 lbs. calcium carbonate equivalent per ton	

## GUARANTEED ANALYSIS

### ACTIVE INGREDIENTS:

Dimethylamine salt of 2, 4-Dichlorophenoxyacetic acid* . . . . .	0.72%
Butyl ether esters of silvex 2-(2, 4, 5-Trichlorophenoxy) propionic acid** . . . . .	0.60%
Technical Chlordane*** . . . . .	1.25%

### INERT INGREDIENTS

Relative to Herbicides . . . . .	97.43%
Relative to Insecticide . . . . .	98.68%
Relative to Insecticide . . . . .	98.75%

\*Equivalent to 0.60% 2, 4-Dichlorophenoxyacetic acid

\*\*Equivalent to 0.40% 2-(2, 4, 5-Trichlorophenoxy) propionic acid (silvex)

\*\*\*Equivalent to 0.75% DDT, 4, 7-Medianoctahydrodane and 0.50% related compounds.

### FERTILIZER

Total Nitrogen (N) . . . . .	10%
7% Ammoniacal Nitrogen	
3% Nitrate Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	6%
From Superphosphate	
Soluble Potash (K <sub>2</sub> O) . . . . .	4%
From Muriate of Potash	
Potential acidity 900 lbs. calcium carbonate equivalent per ton	

## INSTRUCTIONS

This product is formulated to provide a complete feeding for grasses while controlling broadleaf weeds and insects. 1. Kills most broadleaf weeds such as dandelion, plantain, pigweed and similar broadleaf weeds. 2. Controls common lawn insects including ants, white grubs, chinch bugs, wire worms and sod webworms. 3. Feeds lawn for lasting beauty. 4. Apply early Spring, Summer and Fall. For best results apply when weeds are vigorously growing.

1. Spread contents of this bag over 5000 sq ft.
2. Apply when grass is wet from sprinkling or dew.
3. Apply when temperature is 70° F. or higher.
4. Do not mow lawn for 24 hours after application.
5. Do not apply to any other lawn or to surrounding plants.
6. If the product is applied to a lawn and material is watered into soil at a rate of 1/2 inch.

### SPREADER SETTINGS

Spreader	1/2
Settings	1/2
Rate of application	1/2
Cycle time	1/2
Flow rate	1/2

*Handwritten signature*

5772-18

## CAUTION

Avoid inhalation, contact with eyes, nose, mouth, or skin. Wash thoroughly with soap and water after handling. Do not apply to lawns, playgrounds, or other areas where children play.

7% Ammoniacal Nitrogen	
3% Nitrate Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	6%
From Superphosphate	
Soluble Potash (K <sub>2</sub> O)	4%
From Muriate of Potash	
Potential acidity 900 lbs. calcium carbonate equivalent per ton	

### INSTRUCTIONS

This product is formulated to provide a complete feeding for grasses while controlling broadleaf weeds and insects. 1. Kills most broadleaf weeds such as dandelion, plantain, clover and similar broadleaf weeds. 2. Controls common lawn insects including ants, white grubs, chinch bugs, wire worms and sod webworms. 3. Feeds lawn for lasting beauty. 4. Apply in Spring, Summer and Fall. For best results apply when weeds are vigorously growing.

1. Spread contents of this bag over 5000 sq ft.
2. Apply when grass is wet from sprinkling or dew.
3. Apply when temperature is 70° F. or higher.
4. Do not fertilize or mow lawn and do not water lawn for 24 hours after application.
5. Do not apply to lawn areas with existing disease.
6. Do not apply to lawn areas with existing insect damage.
7. Do not apply to lawn areas with existing fungal diseases.
8. Do not apply to lawn areas with existing nematode infestation.

### SPREADER SETTINGS

Spread	1	<i>Klein, 1967</i>
Setting	6	
Control	6	
Check	5	
Control	5	

3172-18

### CAUTION

Avoid inhalation of dusts. Avoid contact with eyes. Avoid contact with skin. Avoid contact with clothing. Avoid contact with children and pets. Avoid contact with water. Avoid contact with food. Avoid contact with feed. Avoid contact with machinery. Avoid contact with electrical equipment. Avoid contact with open flames. Avoid contact with heat. Avoid contact with fire. Avoid contact with explosives. Avoid contact with oxidizing agents. Avoid contact with acids. Avoid contact with alkalis. Avoid contact with toxic substances. Avoid contact with irritants. Avoid contact with sensitizers. Avoid contact with carcinogens. Avoid contact with mutagens. Avoid contact with teratogens. Avoid contact with reproductive toxins. Avoid contact with developmental toxins. Avoid contact with neurotoxins. Avoid contact with immunotoxicants. Avoid contact with immunosuppressants. Avoid contact with immunomodulators. Avoid contact with immunoenhancers. Avoid contact with immunoinhibitors. Avoid contact with immunostimulants. Avoid contact with immunosuppressants. Avoid contact with immunomodulators. Avoid contact with immunoenhancers. Avoid contact with immunoinhibitors. Avoid contact with immunostimulants.

**EARL MAY SEED & NURSERY CO.**  
**Shenandoah, Iowa 51601**