

# Notice

The manufacturer and the seller assume no liability for any injury or damage that may result from the use of this product in any manner other than that specified in the directions. The user is advised to read the directions carefully and to use the product only in accordance with the directions. The user is also advised to use the product only on the surfaces and in the manner specified in the directions. The user is further advised to use the product only on the surfaces and in the manner specified in the directions. The user is also advised to use the product only on the surfaces and in the manner specified in the directions.

# Cautions

Avoid contact with face and hands. If accidentally in contact with face or hands, wash immediately with soap and water. Do not use on children's toys. Do not use on food surfaces. Do not use on clothing. Do not use on painted surfaces. Do not use on electrical equipment. Do not use on electrical equipment. Do not use on electrical equipment.

DO NOT STORE WHERE CHILDREN CAN GET AT IT

MIXING TABLE AND DIRECTIONS

Amount of Spray Desired	Decrease	Increase
1 Quart	2 Teaspoons	2 Teaspoons
1 Gallon	2 Pints	2 Pints
5 Gallons	10 Pints	10 Pints

Table of Equivalents:  
 1 Teaspoon = 1/6 Pint  
 1 Pint = 1/8 Gallon  
 1 Quart = 1/4 Gallon  
 1 Gallon = 1 Gallon

SOLD FOR MERCHANTABILITY



FINE SULPHUR SPRAY

FOR ALL TYPES OF PLANTS, FRUIT AND VEGETABLES

# NOTICE

Table of Equivalents:  
 1 Teaspoon = 1/6 Pint  
 1 Pint = 1/8 Gallon  
 1 Quart = 1/4 Gallon  
 1 Gallon = 1 Gallon

SOLD FOR MERCHANTABILITY

# Directions For Use

(See Mixing Table For Proper Dilutions)

**For Fallage Strengths:** Roses: For control of Black Spot, Powdery Mildew, and Red Spider, apply at 10 to 15 day intervals during season.

**For control of Powdery Mildew and Anthracnose:** apply when shoots are 6 to 8 inches long and again after bloom. Excessive fallage may be somewhat damaged.

**For control of Brown Rot and Scab prevention:** make application at petal fall stage. Do not use on Apricots, Peaches, and Gooseberries; For control of Powdery Mildew, apply before and after blooming.

**For Spring Strengths:** (As growth starts). Roses: For control of Black Spot, Powdery Mildew, and Red Spider, apply at 10 to 15 day intervals during season.

**For control of Powdery Mildew and Anthracnose:** apply when shoots are 6 to 8 inches long and again after bloom. Excessive fallage may be somewhat damaged.

**For control of Brown Rot and Scab prevention:** make application at petal fall stage. Do not use on Apricots, Peaches, and Gooseberries; For control of Powdery Mildew, apply before and after blooming.

**For control of Peach Leaf Curl, San Jose Scale, Brown Rot, and Red Spider:** apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

**For control of San Jose Scale, Powdery Mildew, and Anthracnose:** apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

**For control of Peach Leaf Curl, San Jose Scale, Brown Rot, and Red Spider:** apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

**For control of Peach Leaf Curl, San Jose Scale, Brown Rot, and Red Spider:** apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

# Notice

The manufacturer and the seller assume no liability for any injury or damage that may result from the use of this product in any manner other than that specified in the directions. The user is advised to read the directions carefully and to use the product only in accordance with the directions. The user is also advised to use the product only on the surfaces and in the manner specified in the directions. The user is further advised to use the product only on the surfaces and in the manner specified in the directions.

# Cautions

Avoid contact with face and hands. If accidentally in contact with face or hands, wash immediately with soap and water. Do not use on children's toys. Do not use on food surfaces. Do not use on clothing. Do not use on painted surfaces. Do not use on electrical equipment. Do not use on electrical equipment. Do not use on electrical equipment.

DO NOT STORE WHERE CHILDREN CAN GET AT IT

MIXING TABLE AND DIRECTIONS

Amount of Spray Desired	Decrease	Increase
1 Quart	2 Teaspoons	2 Teaspoons
1 Gallon	2 Pints	2 Pints
5 Gallons	10 Pints	10 Pints

Table of Equivalents:  
 1 Teaspoon = 1/6 Pint  
 1 Pint = 1/8 Gallon  
 1 Quart = 1/4 Gallon  
 1 Gallon = 1 Gallon

SOLD FOR MERCHANTABILITY

# Directions For Use

(See Mixing Table For Proper Dilutions)

**For Fallage Strengths:** Roses: For control of Black Spot, Powdery Mildew, and Red Spider, apply at 10 to 15 day intervals during season.

**For control of Powdery Mildew and Anthracnose:** apply when shoots are 6 to 8 inches long and again after bloom. Excessive fallage may be somewhat damaged.

**For control of Brown Rot and Scab prevention:** make application at petal fall stage. Do not use on Apricots, Peaches, and Gooseberries; For control of Powdery Mildew, apply before and after blooming.

**For Spring Strengths:** (As growth starts). Roses: For control of Black Spot, Powdery Mildew, and Red Spider, apply at 10 to 15 day intervals during season.

**For control of Powdery Mildew and Anthracnose:** apply when shoots are 6 to 8 inches long and again after bloom. Excessive fallage may be somewhat damaged.

**For control of Brown Rot and Scab prevention:** make application at petal fall stage. Do not use on Apricots, Peaches, and Gooseberries; For control of Powdery Mildew, apply before and after blooming.

**For control of Peach Leaf Curl, San Jose Scale, Brown Rot, and Red Spider:** apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

**For control of San Jose Scale, Powdery Mildew, and Anthracnose:** apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

**For control of Peach Leaf Curl, San Jose Scale, Brown Rot, and Red Spider:** apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

**For control of Peach Leaf Curl, San Jose Scale, Brown Rot, and Red Spider:** apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.