

**ACCEPTED**

Feb. 5, 1968

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 3772-19

Best copy available

### Notice

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

### Cautions

Avoid contact with face and hands. In case of contact with skin or eyes, wash thoroughly with plenty of water. May cause skin irritation. Do not use an oil spray within three weeks following a Lime Sulphur spray. Do not use on foliage when temperatures exceed 85° F. Do not spray in the heat of the day. Take care spray drift does not contact light painted surfaces as dark stains or spots may result. Rinse all equipment thoroughly with water after use. Keep out of reach of children.

**DO NOT STORE WHERE PRODUCT WILL FREEZE**

#### MIXING TABLE AND DIRECTIONS

Amount of Spray Desired	Dormant	Early Spring	Summer Foliage
1 Quart	2 Tablespoons	4 Tablespoons	1/2 Teaspoon
1 Gallon	1/2 Pint	1/2 Pint	1 Teaspoon
3 Gallons	3 Pints	1 1/2 Pints	3 Teaspoons
	(6 Cups)		

#### Table of Equivalents:

3 tsp. = 1 the.      2 the. = 1 oz.      16 oz. = 1 pint  
10 the. = 1 cup      2 cups = 1 pint

**SOLD FOR INSECTICIDE AND FUNGICIDE PURPOSES**



## LIME SULPHUR SPRAY

Dormant Clean-Up For Shrubs, Roses, Fruit and Shade Trees

#### ACTIVE INGREDIENTS:

Calcium Polysulfide ..... 30.0%  
Calcium Thioisulfate ..... 2.5%

#### INERT INGREDIENTS ..... 68.5%

(Total equivalent Sulfur in solution ..... 24.0%)

### CAUTION

Keep out of the reach of children.  
See other warnings on back panel.

NET CONTENTS ONE GALLON

Packaged For

**Earl May Seed & Nursery Co.**

EMMENDGAN, IOWA

### Directions For Use

(See Mixing Table For Proper Dilutions)

**Summer Foliage Strength:** Roses: For control of Black Spot, Powdery Mildew, Rust and Red Spider, apply at 10 to 15 day intervals on growing season.

**Grapes:** For control of Powdery Mildew and Anthracnose, apply when shoots are 6 to 8 inches long and again after bloom. Existing foliage may be somewhat damaged.

**Fruit Trees:** For control of Brown Rot and Scab prevention, make one application at petal fall stage. Do not use on Apricots, Currants and Gooseberries: For control of Powdery Mildew, apply just before and after blooming.

**Early Spring Strength:** (As growth starts). Roses: For control of Rose and San Jose Scales, Cane Bearers, and Powdery Mildew, apply in Fall after leaves drop and again in Spring before the buds swell.

**Blackberries and Raspberries:** For control of Red Berry Mite, Anthracnose, Powdery Mildew, and Rust, make one application when canes are 8 to 12 inches tall. Repeat in Fall for Red Berry Mite.

**Dormant Strength:** Apples, Pears, Plums and Cherries: For San Jose Scale control, use before buds swell in Spring. For Pear Leaf Blight Mite, delay until buds start to swell as mites overwinter in the buds.

**Peaches:** For control of Peach Leaf Curl, San Jose Scale, Brown Rot and Twig Borer, apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

**Shrubs and Trees:** For control of San Jose Scale, Powdery Mildew, Canker, Anthracnose, Maple Gall, Leaf Blotch, and Olive Scale, apply once in early Spring before buds swell. Do not apply to Prunus, Eucalyptus, or Rhododendrons, or allow spray to drift on these susceptible species.

U.S. PAT. 2,772,119

G-1169A-J

3772-19

02/05/68

111