

ACCEPTED

AUG 20 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 3772-19

MAYWAY

**LIME SULPHUR
SPRAY**

Dormant Clean-Up For Shrubs, Roses, Fruit and Shade Trees

ACTIVE INGREDIENTS:	
Calcium Polysulfide	30.0%
Calcium Thiosulfate	1.5%
INERT INGREDIENTS	
(Total equivalent Sulfur in solution	34.0%)

CAUTION


Keep out of the reach of children.
See other warnings on back panel.

NET CONTENTS ONE GALLON

Packaged For

Earl May Seed & Nursery Co.

SHEMANDOAN, IOWA



Registrant: Earl May Seed & Nursery Co., Shenandoah, Iowa 51601
Brand Name: MayWay Lime Sulphur Spray 3772-19
Correspondence to: Imperial Inc., P.O.Box 423, Shenandoah, Iowa 51601

Notice

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Cautions

Avoid contact with face and hands. In case of contact with skin or eyes, wash thoroughly with plenty of water. May cause skin irritation. Do not use as oil spray within three weeks following a Lime Sulphur spray. Do not use on foliage when temperatures exceed 85° F. Do not spray in the heat of the day. Take care spray drift does not contact light painted surfaces as dark stains or spots may result. Rinse all equipment thoroughly with water after use. Keep out of reach of children.

DO NOT STORE WHERE PRODUCT WILL FREEZE

MIXING TABLE AND DIRECTIONS

Amount of Spray Desired	Dormant	Early Spring	Summer Foliage
1 Quart	2 Tablespoons	4 Tablespoons	1/2 Cup
1 Gallon	1 Pint	2 Tablespoons	1/2 Cup
3 Gallons	3 Pints (6 Cups)	1/2 Pint	1/2 Tablespoons

Table of Equivalents:
3 tsp = 1 tbs.
16 tbs = 1 cup
2 tbs = 1 oz.
2 cups = 1 pint
16 oz = 1 pint

SOLD FOR INSECTICIDE AND FUNGICIDE PURPOSES

ACCEPTED
Aug 20, 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 3772-19

Directions For Use

Summer Foliage Strength: For control of Black Spot, Powdery Mildew, Rust and Red Spider, apply at 10 to 15 day intervals in growing season.

Grapes: For control of Powdery Mildew and Anthracnose, apply when shoots are 5 to 8 inches long and again after bloom. Rust on foliage may be somewhat delayed.

Peaches: For control of Berry Rot and Scab prevention, make one application at petal fall stage. Do not use on Apricots. **Currents and Gooseberries:** For control of Powdery Mildew, apply just before and after blooming.

Early Spring Strength: (As growth starts). **Roses:** For control of Rose and San Jose Scales, Case Bearers, and Powdery Mildew, apply in Fall after leaves drop and again in Spring before the buds swell.

Raspberries and Blackberries: For control of Red Berry Mites, Anthracnose, Powdery Mildew, and Rust, make one application when new canes are 8 to 12 inches tall. Repeat in Fall for Red Berry Mites.

Dormant Strength: Apples, Pears, Plums and Cherries: For San Jose Scale control, use before buds swell in Spring. For Pear Leaf Blister Mites, delay until buds start to swell as mites overwinter in the bud.

Peaches: For control of Peach Leaf Curl, San Jose Scale, Brown Rot and Twig Borer, apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

Shade Trees and Shrubs: For control of San Jose Scale, Powdery Mildew, Canker, Anthracnose, Maple Gall, Leaf Blotch, and Olive Scale, apply once in early Spring before buds swell. Do not apply to Evergreens, Euonymus, or Rhododendrons, or allow spray to drift on these susceptible species.