

Notice

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Cautions

Avoid contact with face and hands. In case of contact with skin or eyes, wash thoroughly with plenty of water. May cause skin irritation. Keep out of the reach of children.

Do not use an oil spray within three weeks following a Lime Sulphur spray. Do not use on foliage when temperatures exceed 85° F. Do not spray in the heat of the day. Take care spray drift does not contact light painted surfaces as dark stains or spots may result. Rinse all equipment thoroughly with water after use.

Do not reuse empty container. Destroy by burning with waste or burning. Stay away from smoke or fumes.

DO NOT STORE WHERE PRODUCT WILL FREEZE

MIXING TABLE AND DIRECTIONS

Amount of Spray Desired	Dormant	Early Spring	Summer Foliage
1 Quart	8 Tablespoons	4 Tablespoons	1/2 Teaspoon
1 Gallon	1 Pint	5 Tablespoons	3 Teaspoons
3 Gallons	3 Pints (6 Cups)	1/2 Pint	3 Tablespoons

Table of Equivalents:

3 tsp. = 1 tbs. 2 tbs. = 1 oz. 16 oz. = 1 pint
16 tbs. = 1 cup 2 cups = 1 pint


SOLD FOR INSECTICIDE AND FUNGICIDE PURPOSES

PACKAGED FOR
EARL MAY SEED & NURSERY CO.
SHENANDOAH, IOWA



LIME SULPHUR SPRAY

**DORMANT CLEAN-UP FOR SHRUBS, ROSES,
FRUIT AND SHADE TREES**



Summer Fe
dery Milder
in growing

Grates: Fe
when shoot
ing foliage

Peaches: I
one applic
Currants a
just before

Early Sprk
of Rose an
apply in
buds swell

Raspberries
Anthracnos
when new
Berry Mite

Dormant S
Jose Scale
Blister Mit
in the bud

Peaches: F
Rot and T
before bud

Shade Tree
Mildew, Ca
Scale, appl
to Evergre
drift on th

EPA Reg. I

EA

UNDER THE FEDERAL INSECTICIDE
AND FUNGICIDE ACT

07.18.1971

07.18.1971

Notice

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Cautions

Avoid contact with face and hands. In case of contact with skin or eyes, wash thoroughly with plenty of water. May cause skin irritation. Keep out of the reach of children.

Do not use an oil spray within three weeks following a Lime Sulphur spray. Do not use on foliage when temperatures exceed 80° F. Do not spray in the heat of the day. Take care spray drift does not contact light painted surfaces as dark stains or spots may result. Rinse all equipment thoroughly with water after use.

Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

DO NOT STORE WHERE PRODUCT WILL FREEZE

MIXING TABLE AND DIRECTIONS


Amount of Spray Desired	Dormant	Early Spring	Summer Foliage
1 Quart	8 Tablespoons	4 Tablespoons	1/2 Teaspoon
1 Gallon	1 Pint	5 Tablespoons	3 Teaspoons
3 Gallons	3 Pints (6 Cups)	1/2 Pint	3 Tablespoons

Table of Equivalents:

3 tsp. = 1 tbs. 2 tbs. = 1 oz. 16 oz. = 1 pint
16 tbs. = 1 cup 2 cups = 1 pint

SOLD FOR INSECTICIDE AND FUNGICIDE PURPOSES

PACKAGED FOR
EARL MAY SEED & NURSERY CO.
SHENANDOAH, IOWA



LIME SULPHUR SPRAY

**DORMANT CLEAN-UP FOR SHRUBS, ROSES,
FRUIT AND SHADE TREES**

ACTIVE INGREDIENTS:

Calcium Polysulfide 30.0%
Calcium Thiosulfate 1.5%
INERT INGREDIENTS 68.5%
(Total equivalent Sulphur in solution 24.0%)

CAUTION

**Keep out of the reach of children.
See other warnings on back panel.**

NET CONTENTS ONE GALLON

Summer Foliage Mildew in growing

Grapes: Foliage when shooting foliage

Peaches: Foliage one application just before

Early Spring of Rose when buds swell

Raspberries: Anthracnose when new Berry Mite

Dormant Scale Blister Mite in the bud

Peaches: Foliage Rot and Typhlodium before bud

Shade Tree Mildew, Scale, applied to Evergreen drift on the


EPA Reg. No. 1

EA



LIME SULPHUR SPRAY

DORMANT CLEAN-UP FOR SHRUBS, ROSES,
FRUIT AND SHADE TREES



Directions For Use

(See Mixing Table For Proper Dilutions)

Summer Foliage Strength: Roses: For control of Black Spot, Powdery Mildew, Rust and Red Spider, apply at 10 to 15 day intervals in growing season.

Grapes: For control of Powdery Mildew and Anthracose, apply when shoots are 6 to 8 inches long and again after bloom. Existing foliage may be somewhat damaged.

Peaches: For control of Brown Rot and Scab prevention, make one application at petal fall stage. Do not use on Apricots, Currants and Gooseberries. For control of Powdery Mildew, apply just before and after blooming.

Early Spring Strength: (As growth starts). Roses: For control of Rose and San Jose Scales, Case Bearers, and Powdery Mildew, apply in Fall after leaves drop and again in Spring before the buds swell.

Raspberries and Blackberries: For control of Red Berry Mites, Anthracose, Powdery Mildew, and Rust, make one application when new canes are 8 to 12 inches tall. Repeat in Fall for Red Berry Mites.

Dormant Strength: Apples, Pears, Plums and Cherries: For San Jose Scale control, use before buds swell in Spring. For Pear Leaf Blister Mites, delay until buds start to swell as mites overwinter in the bud.

Peaches: For control of Peach Leaf Curl, San Jose Scale, Brown Rot and Twig Borer, apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.

Shade Trees and Shrubs: For control of San Jose Scale, Powdery Mildew, Canker, Anthracose, Maple Gall, Leaf Blotch, and Olive Scale, apply once in early Spring before buds swell. Do not apply to Evergreens, Euonymus, or Rhododendrons, or allow spray to drift on these susceptible species.

EFA Reg. No. 3772-19

K-8265A-G

PACKAGED FOR
EARL MAY SEED & NURSERY CO.
SHENANDOAH, IOWA