

PM 21

3772-19

10/3/97

Pg 176



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OCT 3 1997

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Anne Montalone
Registration Specialist
Bonide Products, Inc.
2 Wurz Avenue
Yorkville, NY 13495

Subject: Labeling Amendment to add "Homeowner Use Only"
Earl May Lime Sulphur Spray
EPA Reg. No. 3772-19
Your Letter Dated July 16, 1996

Dear Ms. Montalone:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable with the following conditions:

1. Delete the Spanish precautionary statement and replace with the following statements:

"Si usted no entienda la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)"

2. In the **PRECAUTIONARY STATEMENTS** section, add the following statements:

"Do not breath dust, vapor or spray mist or get in eyes, on skin or clothing. Wear coveralls over long-sleeved shirt and long pants, socks, shoes, waterproof gloves and use protective eyewear such as goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse."

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release of the product for shipment constitutes acceptance of these conditions. Please submit one copy of your final printed labeling before you release the product for shipment. A stamped copy is enclosed for your records.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


If you have any questions, then please contact Terri Stowe of my staff at (703) 305-6117.

Sincerely,

Mary L. Waller

Mary L. Waller
Acting Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure



LIME SULPHUR SPRAY

Insecticide and Fungicide For Fruit Trees, Shade Trees, Shrubs, Roses and Berries

ACTIVE INGREDIENT	
Calcium Polysulfide.....	50.0%
INERT INGREDIENTS.....	50.0%
TOTAL.....	100.0%
(Total Equivalent sulphur in solution 24.9%)	

KEEP OUT OF REACH OF CHILDREN
PELIGRO — DANGER

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

EPA REG. NO. 3772-19
EPA EST. NO. 4-NY-1

SEE SIDE AND
BACK PANELS
FOR ADDITIONAL
PRECAUTIONARY
STATEMENTS

NET CONTENTS
Manufactured for:
EARL MAY SEED & NURSERY L.P.
Shenandoah, Iowa 51603

ACCEPTED
with COMMENTS
In EPA Letter Dated

OCT 2 1967
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

3772-19

4/6

Lime Sulphur Spray

Insecticide and Fungicide for Fruit Trees,
Shade Trees, Shrubs, Roses and Berries

EPA Est. No. 4-NY-1 (1), 769-GA-1 (2), 55179-IN-1 (3), 44616-MO-1 (4).
EPA Reg. No. 3772-19
The last digit in the lot number indicates EPA Establishment location.



SATISFACTION GUARANTEE:
If, for any reason you are not completely satisfied with the results after applying this product, you are entitled to product replacement or your money back simply by presenting evidence of purchase to your local Earl May Nursery & Garden Center.

Manufactured for
**EARL MAY SEED
& NURSERY**
Shenandoah, IA 51603

1 Quart

Fix-a-Form PAT. 448822/456043
MOSS PRINTING, CHICAGO, IL 60645

DORMANT SPRAY

Apples, Pears, Plums and Cherries: For San Jose Scale, use before buds swell in Spring. For Pear Leaf Blister Mites, delay until bud swell to treat overwintering mites in buds.

Peaches: For control of Peach Leaf Curl, San Jose Scale, Twig Borer and as an aid in the control of Brown Rot Blossom Blight; apply once in late fall, repeat in spring just before buds swell. Full season control of Brown Rot Blossom Blight will require additional application of a fungicide.

Shade Trees and Shrubs: For control of San Jose Scale, Noctria Canker, Maple Gall, Leaf Blotch and Olive Scale, and as an aid in the control of Powdery Mildew and Anthracnose; apply once in early Spring before buds swell. Do not apply to Evergreens, Euonymus or Rhododendrons or allow spray to drift on these susceptible species. Full season control of Powdery Mildew and Anthracnose will require additional application of a fungicide.

EARLY SPRING SPRAYING (AS GROWTH STARTS)

Roses: For control of Rose and San Jose Scale, Case Boarers and Powdery Mildew; apply in Spring before buds swell and again in fall after leaves drop.

Raspberries and Blackberries: For control of Red Berry Mites, and as an aid in the control of Anthracnose, Powdery Mildew, and Rust; make one application when canes are 8 to 12 inches high. Repeat in Fall for Red Berry Mites. Full season control of Anthracnose, Powdery Mildew, and Rust will require additional application of a fungicide.

SUMMER FOLIAGE SPRAYING

Roses: For control of Black Spot, Powdery Mildew, Rust and Red Spider;

apply at 10 to 15 day intervals in growing season.

Grapes: As an aid in the control of Powdery Mildew and Anthracnose, apply when shoots are 6 to 8 inches long and again after bloom. Existing foliage may be somewhat damaged. Additional applications of a fungicide are necessary for full seasonal control.

Peaches: As an aid in the control of Brown Rot Blossom Blight and to help prevent Scab; make an application at petal fall stage. Do not use on Apricots, Currants or Gooseberries. For control of Powdery Mildew; apply just before and after blooming. Additional application of a fungicide may be necessary for full season control.

Amount of Spray Desired	Dormant	Early Spring	Summer Foliage
1 Quart	8 Tablespoons	4 Tablespoons	3/4 Teaspoon
1 Gallon	1 Pint	16 Tablespoons	3 Teaspoons
3 Gallons	3 Pints (6 cups)	1 1/2 pint	3 Tablespoons

Table of equivalents:

3 tsp. = 1 Tbs. 2 Tbs. = 1 oz. 16 oz. = 1 pint
16 Tbs. = 1 cup 2 cups = 1 pint

COMBINATION WINTER SPRAY

Fruit Trees, Roses and Deciduous Hedge Plants, Shade Trees and Shrubs (Dormant and Deciduous Only)

San Jose Scale, Rose Scale, Oyster Shell Scale, Brown Apricot Scale, Black

Scale, Overwintering Insect Eggs, many Overwintering Fungus Spores or Plant Disease.

FOR BEST RESULTS: Use 1-1/2 pints Earl May Lime Sulphur Spray and 1/2 pint Earl May All Seasons Horticultural & Dormant Spray Oil in 6 gallons of water (7-1/2 tablespoonsfuls Earl May Lime Sulphur Spray and 2-1/2 tablespoonfuls Earl May All Seasons Horticultural & Dormant Spray Oil per gallon). Keep agitated during spraying.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER

Fatal if swallowed, inhaled or absorbed through skin. This product is corrosive to flesh because of its caustic alkaline nature and it causes irreversible eye damage and skin burns.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink promptly a large quantity of milk, egg white, or gelatin solution or if these are not available, large quantities of water. Get medical attention. Do not induce vomiting.

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush with plenty of water. Get medical attention.

ENVIRONMENTAL HAZARD:
 Do not apply directly to water. Do not contaminate water used for domestic or irrigation purposes.

DIRECTIONS FOR USE

DD "Homeowners Use Only"

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

STORAGE AND DISPOSAL

Storage: Store in original container, in a cool, dry, secure area, away from fertilizer, food or feed. Keep container closed. Do not store where product will freeze.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty container. Wrap container and put in trash.

Do not use on evergreens, euonymus, rhododendrons, or allow spray to drift on to these susceptible species.

Do not use an oil spray within three weeks after a Lime Sulphur application. Do not spray in heat of day or when temperatures exceed 85°F. Do not let spray contact painted surfaces as dark stains or spots may result.