

\$1.98

10% CHLORDANE GRANULES



CAUTION
KEEP OUT OF THE REACH OF CHILDREN.
See other warnings on back panel.

ACTIVE INGREDIENTS:
Technical Chlordane* 10%
INERT INGREDIENTS: 90%
*Equivalent to 6% Octachloro-4, 7-methanotetrahydroindane
and 4% related compounds


NET WEIGHT 3 POUNDS

FOR SALE BY NURSERY

MAY WAY

**10%
CHLORDANE GRANULES**

FOR
SOIL
INSECT
CONTROL



FOR
SOIL
INSECT
CONTROL

MANUFACTURED FOR

EARL MAY SEED & NURSERY CO.

SHENANDOAH, IOWA

Feb. 23, 1971

3772 22

BACK PANEL

CORRECTED LABELING

EARL MAY 10% CHLORDANE GRANULES

GENERAL DIRECTIONS

10% Chlordane Granules should be thoroughly mixed into the soil prior to planting or transplanting in the Spring.

BROADCAST APPLICATIONS: Apply the proper amount of 10% Chlordane Granules to the soil. (See chart). After application, stir or mix insecticide in the soil as uniformly as possible.

ROW OR BAND TREATMENT (Use at preplanting, planting or transplanting time only): For the protection of row crops, apply 10% Chlordane Granules in a band 6 to 8 inches wide. (See chart). After application, stir or mix insecticide in the soil as uniformly as possible. Do not use band treatment on potatoes grown by ridging cultivation practice. It should be noted that band treatments are effective for one growing season only.

INCREASE DOSAGE 50% ON PEAT OR MUCK SOILS

Crop	Insects	VEGETABLES	
		Brdcast	Row or Band (Use at pre-planting, planting, or transplanting time only)
Beans, Black-eyed Peas	White Fringed Beetle	3/4 lb.	---
	Grubs, Wireworms	1/2 lb.	---
	Cutworms	1/2 lb.	---
Rutabagas	Cutworms	1/2 lb.	---
	Wireworms, White Grubs	3/4 lb.	1/4 lb.
Broccoli, Brussels Sprouts, Kohlrabi, Cabbage	Cabbage Maggot	3/4 lb.	---
	Ants, Cutworms, Flea-beetle Larvae, Wireworms	3/4 lb.	---
Cauliflower	Cabbage Maggot	3/4 lb.	---
	White Grubs	3/4 lb.	---
Corn, Sweet and Popcorn	Ants, Corn Rootworm Larvae, Japanese Beetle	1/2 lb.	---
	Grubs, White-Fringed Beetle Larvae	1/2 lb.	---
	Wireworms, Fleabeetle Larvae	3/4 lb.	1/8 to 1/4 lb.

ACCEPTED
Feb. 23, 1971
3772 22

BACK PANEL (cont.)

CORRECTED LABELING

EARL MAY 10% CHLORDANE GRANULES

VEGETABLES			
Crop	Insects	Pounds per 1,000 sq. ft.	
		Brdcast	Row or Band (Use at pre-planting, planting, or transplanting time only)
Peas	Ants, Rootworms, Green June Beetle Grubs	1/2 lb.	---
	Wireworms, White-fringed Beetle Grubs, White grubs	3/4 lb.	---
MAY BE INJURIOUS TO SOME VARIETIES			
Melons, Cucumber, Squash	White Grubs, Wireworms, Cucumber Beetle Larvae	3/4 lb.	1/8 to 1/4 lb.
	Cutworms	1/2 lb.	---
Eggplant	White Grubs, Fleabeetle Larvae	3/4 lb.	---
Peppers	Wireworms	3/4 lb.	1/8 to 1/4 lb.
	Cutworms, Mole Crickets	1/2 lb.	---
Lettuce	Cutworms	1/2 lb.	---
	Wireworms, White Grubs	3/4 lb.	---
DO NOT USE ON GREEN OR SPRING ONIONS			
Onions	Wireworms	3/4 lb.	---
	Onion maggot	1/2 lb.	1/8 to 1/4 lb.
Potatoes	Cutworms	1/2 lb.	---
	Fleabeetle Larvae, Wireworms, White Grubs, Mole Crickets	3/4 lb.	---
	Cutworms	1/2 lb.	---
Sweet Potatoes	Fleabeetle Larvae, Wireworms, White Grubs, White-fringed Beetle Grubs	3/4 lb.	---
	Cutworms	1/2 lb.	---
Tomatoes	Fleabeetle Larvae, Wireworms, White Grubs	3/4 lb.	---

AC
1/6 22 1971
3772-22

BACK PANEL (cont.)

CORRECTED LABELING

EARL MAY 10% CHLORDANE GRANULES

CAUTION: Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

This product is toxic to fish and wildlife. Keep out of any body of water.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ANTIDOTE: External --- In case of spillage on skin, wash promptly with soap and water. Internal --- Emetic of 1 table-spoonful mustard in tumbler of water. Call a physician.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

EARTHWORMS: Apply 1 to 2 lbs. 10% Chlordane Granules per 200 sq. ft. (equal to 20 to 40 pounds actual Chlordane per acre). After application, water the insecticide into the soil. This places the insecticide in the most effective control area. Use only on Lawns or Golf Courses as the dosage of actual Chlordane recommended exceeds those accepted on soil to be planted in food or feed crops. Birds feeding on treated areas may be killed.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA Reg. No. 3772-22

1/6 22 1971
3772-22