

...LEAF HOPPER ... PO
 ...FLA BEE
 ...TOMATO HORN
 ...CINER
 ...MARLEQUIN ...
 ...

6% CHLORDANE DUST

CONTENTS OF TUBE

10000

GARDEN HOME LAWN

NOTE

ACTIVE INGREDIENTS:	
Technical Chlordane*	6%
INERT INGREDIENTS	94%
Total	100%

*Equivalent to 3.6% Octachloro-4,7-methanotetrahydroindane and 2.4% related compounds.

CAUTION
KEEP OUT OF THE REACH OF CHILDREN.
 See other warnings on back panel.
NET WEIGHT 1 POUND

DIRECTIONS FOR USE

DIRECTIONS

Apply as directed to control insects by poisoning their food source.

ANTS

IN LAWNS Apply to trails and water troughs and other ant trails. Also apply to areas where ants are nesting.

IN GAR-DENS Apply to trails and water troughs and other ant trails. Also apply to areas where ants are nesting.

CAUTIONS Avoid contact with children and pets. Do not apply to areas where children and pets play.

GARDEN PESTS

IN HOMES Apply to cracks and crevices, under baseboards, and behind doors and windows.

IN HOMES Apply to cracks and crevices, under baseboards, and behind doors and windows.

LAWNS

To control ants, chiggers, chinch bugs, etc., apply this dust at the rate of 1 pound per 1,000 sq. ft. Water material down into soil thoroughly. Keep children and pets off the treated areas until the insecticide has been washed into the soil and treated areas have dried completely.

SOIL INSECTS

LAWNS. To control **WHITE GRUBS** apply at the rate of 1 pound per 400 square feet. For **JAPANESE BEETLE GRUBS** use 1 pound per 250 square feet. Wash material into soil with ample quantities of water to insure deep penetration. **SOD WEBWORM (LAWN MOTHS).** Use 1 pound per 400 square feet; water thoroughly before treatment and do not water again for several days. Remains effective entire season. **GARDENS.** To control wireworms, white grubs, Japanese beetle grubs attacking potatoes, corn, cucumbers, tomatoes, peas, and beans, apply at the rate of 1 pound per 500 square feet before planting and work into the top 2 or 3 in. of soil. Remains effective entire season.

IN HOMES

Apply as directed below. Repeat applications as needed to maintain effective control.

ANTS. See directions under **ANTS.** **ROACHES, WATERBUGS, SILVERFISH.** Dust into infested cracks, under sinks, behind baseboards and around and nesting places of these insects.

SILVERFISH, also treat shelving and bookcases. **BOX ELDER BUGS.** Apply around doorsills, window frames, or wherever these pests enter premises.

Apply to outside areas and places where these insects congregate. **CRICKETS, SPIDERS, CENTIPEDES.** Apply dust to dark places and crevices in basements, closets, and behind baseboards.

CARPET BEETLES. Apply to area of the floor, baseboards, and closet shelves and work the dust into cracks where the insects may be found. Treat the underside of carpet.

GARDEN PESTS

ON FLOWERS. To control **ANTS, ARMYWORMS, CUCUMBER BEETLES, FLEA BEETLES, GRASSHOPPERS, CLIMBING CUTWORMS, LEAF MINERS, TARNISHED PLANT BUGS, THRIPS** on gladiolus, gladiolus, roses, zinnias, other flowers, ornamentals and shrubs, apply to plants.

Apply to flowers and in crevices of plants. Repeat applications as needed.

ON VEGETABLES. To control **COLORADO POTATO BEETLE, POTATO LEAFHOPPER, POTATO FLEA BEETLE, ON BEANS, FLEA BEETLE, GRASSHOPPERS, TOMATO HORNWORMS, ON TOMATOES: LEAF MINERS on peppers, CABBAGE WORMS, HARLEQUIN BUGS, CABBAGE LOOPERS on cabbage, SPITTLE BUGS, STRAWBERRY CROWN-BORER on strawberries, BEAN LEAF ROLLERS, STINK-BUG, on beans; apply thoroughly to plants cover both upper and under sides of foliage. Repeat as needed. To aid in the control of **ONION MAGGOTS** on onions, apply the dust around the base of plants as soon as they are set in the ground. The insecticide should be washed or stirred in soil. Do not apply to spring or green onions.**

SPECIAL CAUTIONS. Do not use on cucurbits or on cabbage/after heads or other edible portions begin to form; or tomatoes, peppers, strawberries after fruit begin to form; nor on green beans or snap beans after pods begin to form. Do not use on leafy vegetables such as lettuce, spinach. Remove residues remaining at harvest by washing, wiping, or other effective means. Do not apply to food crops within 30 days of harvest, nor to forage crops to be fed to dairy animals or animals being finished for slaughter.

TREATMENT OF BULBS

Before storing bulbs and for treatment just before planting to control trips on Dahlias, Hyacinths, Tulips, Gladiolus, and most flowering bulbs, place new bulbs in a strong paper bag, dust a table with 1/2 lb. Chlor-dane Dust on the bulbs, closing the bag and shake vigorously.

CAUTION

Irritant if swallowed. Avoid inhalation of dust. Avoid contact with skin. Wash hands after using. Avoid contamination of food, foodstuffs, cooking utensils. Keep out of reach of children. This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of **MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE**, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Packed by weight, not in volume. Product may set in shipment.


EPA Reg. No. 3772-25 K-8266A-G

Manufactured for **EARL MAY SEED & NURSERY CO. SHENANDOAH, IOWA**



6% CHLORDANE DUST

for the control of insects infesting the



GARDEN - HOME - LAWN

ACCEPTED
 Nov 16, 1971
 5772-25