OCT 01 1991

Mr. Rocky Steinbrink
May Seed & Nursery L.P.
208 North Elm Street
P.C. Box 423
Shenandoah, IA 51603

Dear Mr. Steinbrin::

Subject: Earl May Garden Rotenone Dust
EPA Registration No. 3772-32
Your Submission of a Drait Label
With the "Generic Data Exemption
Statement" Dated June 1, 1989
Responding to the Rotenone
Registration Standard

The draft label you submitted is acceptable. A stamped copy is enclosed for your file.

The latest Confidential Statement of Formula (CSF) on file for the product does not have certified limits. Submit a revised CSF within 60 days of the date stamped on this letter.

Sincerely yours,

Phil Hutton

Product Manager (10)

Insecticide-Rodenticide Branch Registration Division (H7505C)

Enclosure

50580:I:Schroeder:L18-31:KEVRIC:06/20/91:DD:WO:CL

CONCURRENCES								
SYMBOL								
SURNAME								
DATE					***************************************	***************		

EPA Form 1320-1 (12-70)

OFFICIAL FILE COPY

(Front Panel)

EARL

# GARDEN ROTENONE DUST

For Vegetables & Flowers

ACTIVE INGREDIENTS:
Rotenone
Other Associated Resins
INERT INGREDIENTS
Total
1.66%
3.34%
95.00%


KEEP OUT OF REACH OF CHILDREN

# CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT\_\_\_\_





(Side Panels)

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harnful if swallowed. Avoid breathing dust. Avoid contact with skin.

#### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic orginisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply to vegetable crops within one day of harvest.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried or dust have settled.

STORAGE: Store in original container, in a cool, dry, secure area away from fertilizer, food or feed. Keep container closed.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

## HIGHT CRAWLERS

NIGHT CRAVLERS -- sometimes called red worms or fish worms -- can cause considerable damage to lawns in some areas. The unsightly mounds of soil thrown up by night crawlers creates a rough uneven lawn. This condition usually occurs during early moist spring weather when night crawlers are most active near the surface.

Apply EARL MAY GARDEN ROTENONE DUST at the rate of 1 lb. to 250 square feet of lawn. Sift from shaker top can or use a small nand duster. Water the material into the ground. For best quick results make application just before a spring shower when night crawlers are coming to the top for feeding.

#### PETS

Effective against fleas, lice and ticks on ponies, dogs and cats. Sift the Rotenone Dust into the hair and rub in by hand. Do not use on puppies or kittens less than 4 weeks old.

EARL MAY ROTENONE DUST gives good control of a wide variety of both sucking and chewing insects infesting vegetable and flower gardens. Non-woolly aphids (plant lice), bean leaf beetles, flea beetles, imported cabbage worm, sawfly larva, spittlebug, asparagus beetle.

BEST AVAILABLE CUPY

EARL MAY ROTENONE DUST may be used with safety on flowers, ornamental plants and cabbage, cauliflower, lettuce, beans, tomatoes, peas, radishes, beets, unions, squash, cucumbers, melons and asparagus. Will not injure tender plant foliage. For best results use as a dust applied with small hand duster, or sift lightly from this shaker type can. Dust underneath and top of foliage thoroughly. Apply to both plants and insects. Begin application when plants are well established or when new growth appears. Repeat every 7 to 10 days or as often as necessary, whether insects are present or not to help prevent infestation. Repeat after heavy rains. Dust when air is calm.

No mixing -- ready to use. Dust plants thoroughly, covering all insects to be controlled. May be used as a wet spray. Stir or mix 3 to 4 tablespoonfuls with enough water to make a paste first, then add water to make 1 gallon of spray.

#### NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of NERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Packed by weight, not in volume. Product may settle in shipment.

EPA Reg. No. 3772-32 EPA Est. 407-IA-1 A-72-84-1

MANUFACTURED FOR MAY SEED & NURSERY L.P. SHENAIDOAN, IOWA 51603

