### GLASSES, DISHES, UTENSILS

- 1. Scrape and preflush articles to be washed.
- 2. Wash with recommended Crouch product.
- 3. Rinse in clean water.
- 4. Sanitize with a solution of Crouch Steep (1 oz. per 2 gallons water 200 ppm) by immersing all glassware, dishes and utensils for at least 2 minutes (or for contact time specified by governing sanitary code). Test solution periodically. Discard solution when 5 drops or less of acid testing solution gives red end-point (100 ppm or less).
- 5. Air dry on a rack or drain board.

CONTACT TIME: The laws of some states require that utensils and equipment remain in contact with germicidal solution for 2 minutes. Check your Health Department

MILKSTONE AND WATERSTONE REMOVAL: To remove milkstone or waterstone, use a solution of 1 part Crouch Steep to 2 parts warm water. Dab this solution over the surface and let soak for 2 to 5 minutes, then brush and flush with clear water.

# USE CONCENTRATIONS FOR EVERY TWO GALLONS OF WATER

USE	Dilution Strength	For
1 oz.	200 ppm active* ingredient	Sanitizing by immersion, brushing or circulation.

\*ppm means parts of Crouch Steep active (Dodecyl Benzene Sulfonic Acid) per each million parts of water.

CROUCH STEEP fulfills the criteria of Appendix F, the Grade "A" Pasteurized Milk Ordinance, 1965 Recommendations of the United States Public Health Service.

**Net Contents 1 Gallon** 

# GREGET STEEP A LIQUID ACID SANITIZER CONCENTRATE

## **ACTIVE INGREDIENTS:**

Phosphoric Acid		. 30.00%
Dodecyl Benzene Sulfonic Acid		5.00%
INERT INGREDIENTS:		65.00%
	TOTAL	100.00%

DANGER: KEEP OUT OF REACH OF CHILDREN SEE SIDE PANEL FOR FURTHER PRECAUTIONS

# CROUCH SUPPLY COMPANY, INC.

General Offices: P. O. Box 906 • Fort Worth, Texas 76101

Fort Worth; Houston; San Antonio; El Paso, Texas; and Little Rock, Arkansas.

### RECOMMENDED FOR USE IN DAIRIES, FOOD PROCESSING, BEVERAGE, CHEESE AND ICE CREAM PLANTS ALSO FOR USE ON DAIRY FARMS

CROUCH STEEP Liquid Sanitizer is a concentrate. In case of contact with eyes or skin, flush with plenty of water. If discomfort persists, get medical attention.

DO NOT USE CROUCH STEEP WITH SOAPS OR ALKALINE CLEANERS.

AVOID CONTACT WITH FOOD-STUFFS.

DO NOT USE ON FERROUS OR SILVER METALS.

DIRECTIONS FOR SANITIZING WITH CROUCH STEEP

FOR SANITIZING: Prepare a solution of sufficient volume by adding 1 oz. of Crouch Steep to every 2 gallons of water. Use stainless steel or plastic containers. Sanitize equipment by circulation, immersion or spraying. Allow a 2 minute exposure period or contact time specified by governing sanitary code. Rinse with potable water before reuse.

# IN DAIRY, CHEESE, BEVERAGE, ICE CREAM, FOOD PROCESSING PLANTS

- 1. Clean equipment immediately after use.
- a. Rinse—with clean, tepid water.b. Wash thoroughly with recommended Crouch prod-
- c. Rinse-with clean tepid water.
- d. Air dry—protect from dust and dirt, if possible.
  2. Sanitize immediately before use with a solution of Crouch Steep (1 oz. per 2 gollons water 200 ppm), by soak—circulation or immersion.

# RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD IT.

DANGER: Corrosive: Keep Out of Reach of Children. May be fatal or harmful if swallowed. Do not breathe vapor or fumes. Produces chemical burns. Do not get in eyes, on skin, or on clothing.

ANTIDOTE: If swallowed call a physician immediately. Drink a teaspoonful or more of milk of magnesia or a cup of soapy water followed with milk or raw egg whites.

Eye contact — Wash with water for 15 minutes. Get prompt medical attention.

Skin contact — Wipe off the acid gently, immediately flood the surface with water, using soap freely, then cover with moist magnesia or baking soda.

ACCEPTED

AGAI 11 1973

INDER THE FEDERAL INSECTICIDE
HOW HOLD RODENTICIDE ACT
HOLD HOLD RODENTICIDE
HOLD RODENTICIDE ACT
HOLD RODENT

38/3-14