

3838-37

02/17/1987

1/1

QUAT RINSE

spray at 3 RPM. 5. Fill sprayer reservoir with 1.5 oz. of concentrate to 1 gallon of water to produce 1150 ppm active ingredients. 6. Set sprayer mechanism to deliver 1 gallon of solution. 7. Fog for 15 minutes for an average 2100 cu. ft. room. For different room sizes, vary spray time proportionately to ensure complete wetting of exposed surfaces. 8. Wait 2 hours before entering room after treatment then manually disinfect room surfaces according to recommended procedures. 9. Rinse fogging equipment thoroughly with clear water following use.

COLD DISINFECTION: For disinfection of previously cleaned surgical instrument, barber tools and dental equipment, simply submerge instruments into solution containing 1-1/2 ounces per gallon of water for 10 minutes. Prepare a fresh solution daily or when used solution becomes visibly dirty.

PRECAUTIONARY STATEMENTS

Hazard to Human and Domestic Animals

DANGER—CORROSIVE

Keep out of Reach of Children. Causes severe eye and skin damage. Do not get in eyes, skin or on clothing. Harmful if swallowed. Wear goggles or face shield and rubber gloves when handling. Avoid contamination of food. Remove and wash contaminated clothing.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 min. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

STORAGE & DISPOSAL

- DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

PESTICIDE DISPOSAL — Pesticides that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide.

CONTAINER DISPOSAL - Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers.

QUAT RINSE

DISINFECTANT — DEODORIZER — SANITIZER

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides 5%
n-Alkyl (68% C₁₂, 32% C₁₄)
dimethyl ethylbenzyl ammonium chlorides 5%

INERT INGREDIENTS 90%
100%

KEEP OUT OF REACH OF CHILDREN

DANGER

CAUSES EYE DAMAGE.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON LEFT PANEL.

EPA Reg.No. 3838-37 EPA Est. No. 3838-WI-1

DILUTE CONCENTRATE BEFORE USING

Only for Sale To, Use, and Storage by Service Persons.

Contents: One Gallon

ACCEPTED
3838-37
FEB 17 1987
Under the FIFRA Act, this pesticide is registered under EPA Reg. No. 3838-37. It is restricted to use by service persons only. It is not to be used on food, feed, or animal products.

REGISTERED TRADEMARK
DISTRIBUTORS, INC.

QUAT RINSE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOOD PROCESSING EQUIPMENT: For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce per 4 gallons of water to provide 200 ppm of active quaternaries. Apply with cloth, mop or sponge. Surfaces should remain wet for 1 minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when used solution becomes visibly dirty. For mechanical application, used solution may not be reused for sanitizing applications.

HOSPITALS: For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use concentrate at 3-1/2 ounces per 5 gallons of water. Apply with cloth, mop, or sponge. Treated surfaces must remain wet for 10 minutes. Prepare a fresh solution daily or when used solution becomes visibly dirty. At this level concentrate will be effective against *Pseudomonas aeruginosa* PRD-10. For heavily soiled or contaminated areas, a precleaning step is required.

DAIRIES: To sanitize dairy equipment such as tanks, lines, pails, and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce to 4 gallons of water (200 ppm). Apply with cloth, mop or sponge. Surfaces should remain wet for 1 minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when used solution becomes visibly dirty. For mechanical application, used solution may not be reused for sanitizing applications. Follow recommendations of local Health Board.

HOUSEHOLD: For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Lactar Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce to 4 gallons of water. Disinfect with 3 ounces to 5 gallons of water. Treated surfaces must remain wet for 10 minutes. Prepare a fresh solution daily or when used solution becomes visibly dirty. Kills *Staphylococcus aureus* and *Salmonella choleraesuis*. For heavily soiled or contaminated areas, a precleaning step is recommended.

RESTAURANT AND BAR RINSE: 1. Scrape and prewash utensils and glasses whenever possible. 2. Wash with a good detergent or compatible cleaner (a commercial cleaner or detergent can be recommended). 3. Rinse with clean water. 4. Sanitize in a solution of 1 oz. to 4 gal. of water (200 ppm). Immerse all utensils for at least two minutes or for a contact time specified by governing sanitary code. 5. Place sanitized utensils on a rack or drainboard to air dry.

FOGGING: To minimize the danger of cross contamination as an adjunct to acceptable manual cleaning and disinfecting of room surfaces follow these special directions. 1. Remove all human, animal, and plant life from room. 2. Open closet doors and drawers. 3. Set up 34" revolving platform in center of room. 4. Mount Jet Challenger Sprayer, Model 5100C, or other fogging device delivering equivalent