

EARL MAY GARDEN WEED PREVENTER

GENERAL INFORMATION

Earl May Garden Weed Preventer is a selective herbicide which controls many of the most troublesome weeds in flower gardens, vegetable gardens, and lawns. It is safe for use on all plants listed on the label. This preventive is effective against many of the most common weeds in flower gardens, vegetable gardens, and lawns.

DIRECTIONS

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

RECOMMENDATIONS

FLOWER GARDENS: Apply after establishment or as supplemental application at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds.

Where possible, this should be done in the spring. Weed control is best when proper application is made.

LAWNS: Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE, AND ONIONS:

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

BEANS (dry and snap):

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

POTATOES (whole or cut pieces):

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

TOMATOES:

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

STRAWBERRIES (new plantings):

Apply to the soil in the garden or lawn at the rate of 1 pound per 1,000 sq. ft. Apply to the soil following establishment or setting out of flower beds. Where possible, this should be done in the spring. Weed control is best when proper application is made.

CAUTION

Keep out of the reach of children. Do not use on lawns or in flower gardens.

NOTICE

Do not use on lawns or in flower gardens. Do not use on lawns or in flower gardens.

MANUFACTURED FOR
EARL MAY SEED & NURSERY CO.

SHELANDOAH, IOWA

Dispose of empty container by burning with lid on or burning. Stay away from smoke or fumes.