

**INSTRUCTIONS**  
Application for control of aphids and mites on flowers mix 1 1/2 oz. concentrate per gallon of water. For control of leaf hoppers and thrips on cabbages and clover mite on tomatoes mix 1.0 lb. per acre (4 fl. oz. product per 100 gal.) of water. For control of Japanese beetles on pears use 1 teaspoonful per gallon of water. For control of Mexican bean beetles on beans mix 1 1/2 teaspoonfuls per gallon of water. For control of tent caterpillars, canker worms, certain leaf miners and certain scale crawlers on ornamentals use 2 teaspoonfuls per gallon of water. (2 pints in 100 gallons). Spray when insects appear and repeat when necessary. Do not apply to food crops later than 7 days before harvest. Injury may occur on certain ferns and some species of Crassula. May injure **Bosc pears in Northeast**. May injure McIntosh and Cortland varieties of apples if applied within four weeks of harvest.

**FLEA CONTROL** - To control fleas in kennels, yards, lawns and under houses, dilute 5 ounces in one gallon of water and apply at the rate of one gallon of diluted spray per 1000 square feet. Clean area before treatment. Fresh bedding should be placed in animals quarters following treatment. Repeat as necessary. **DO NOT APPLY TO DOGS.**

**MOSQUITO CONTROL** - For the control of adult mosquitoes, flies and small flying insects in outdoor areas, dilute 1 to 28 with water (4 1/2 fl. oz. per gal.) Apply as fine mist or fog. Three fl. oz. of concentrate will treat 1 acre. Repeat application as necessary. May cause spotting on automobile paint finish. Cars should not be sprayed directly. If accidental exposure occurs, the car should be washed immediately.

**ROACH CONTROL** - For control of roaches, ants, spiders and clover mites in and about homes, institutes and industrial plants, dilute 4 ounces per gallon of water. Apply by means of a coarse spray or paint brush to window frames, base boards, under sinks, drain boards, stoves, and other areas frequented by insects. Repeat as necessary. Do not contaminate food or utensils. Do not use in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. Do not use where milk is stored.

# MALATHION-50 SPRAY

CONTAINS CYTHION®  
PREMIUM GRADE MALATHION

## ACTIVE INGREDIENTS

Malathion\* . . . . . 50.0%

Xylene . . . . . 38%

INERT INGREDIENT . . . . . 12%

Total 100.0%

\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

## CAUTION:

Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Manufactured by:

A.B.C. COMPOUNDING CO., INC.

P.O. BOX 932

ATLANTA, GEORGIA 30301

ACCEPTED

ANT MO  
water. Ma  
beds and  
necessary.

MILLIPE  
around he  
gallon per  
Repeat in  
grown.

HARMFUL  
SKIN. Ma  
is anidota  
with atrop  
clothing.  
with feed  
eyes, get

This prod  
where run  
drift from  
or dispos  
This prod  
on crops.  
Agricultu  
ughly wit  
disposal.

# MALATHION-50 SPRAY

CONTAINS CYTHION®

PREMIUM GRADE MALATHION

## ACTIVE INGREDIENTS

Malathion\* . . . . . 50.0%

Xylene . . . . . 38%

INERT INGREDIENT . . . . . 12%

Total 100.0%

\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

## CAUTION:

Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ANT MOUNDS - Spray hills thoroughly, using 1½ teaspoon per gallon of water. Mix moist earth on the top of the hill. For other small ants in flower beds and lawns, spray lightly in the infested areas. Repeat in 10 - 15 days if necessary.

MILLIPEDES, SPRINGTAILS & SOWBUGS\* in and around greenhouses and around homes. Mix 1 teaspoon in 1 gallon of water and apply at the rate of 1 gallon per 150 sq. ft (10 ft. x 15 ft.) of soil surface where insects congregate. Repeat in 7 - 10 days if necessary. Do not apply to soil where food crops are grown.

## CAUTION

HARMFUL IF INHALED, SWALLOWED OR ABSORBED THROUGH THE SKIN. Malathion is a cholinesterase inhibitor. Note to Physicians: Atropine is antidotal. 2 PAM is also antidotal and may be administered in conjunction with atropine. Avoid breathing spray mist. Avoid contact with skin eyes and clothing. Store away from children and domestic animals. Avoid contact with feed and foodstuffs. In case of contact, wash with soap and water. For eyes, get medical attention.

DO NOT USE, POUR, SPILL, OR STORE  
NEAR HEAT OR OPEN FLAME

## CAUTION

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Wash thoroughly with water and detergent. Wrap in newspaper and place in trash can for disposal.

E.P.A. REG. NO. 3862-47

NET CONTENTS LIQUID GALS.

Manufactured by:

A.B.C. COMPOUNDING CO., INC.

P.O. BOX 932

ATLANTA, GEORGIA 30301

ACCEPTED