

UNDER
 FUNGICIDE
 FOR DOWN
 ED UN... NO. 3862-53

EPA REG NO 3862-53-
EPA EST NO. 3862-GA-1

LEM-O-DIS 19-30

USE DIRECTIONS

Use at 2 ounces per gallon. This product will provide 500 ppm quaternary to the use solution. This is enough for general disinfection, fungicidal activity against pathogenic fungi and deodorizing.

For best results, use **LEM-O-DIS 19-30** with a cloth or mop and apply to walls, floors, table tops, counters and garbage pails.

For use in hospitals, 3 ounces per gallon will be required for disinfection against *Pseudomonas aeruginosa*.

For heavy soil, a pre cleaning step is recommended before using **LEM-O-DIS 19-30**.

Sanitization

When used for sanitization of previously cleaned food processing or food contact items, limit the level of active quaternaries to 200 ppm (4 ounces per 5 gallons of water) and submerge article in sanitizing solution or apply sanitizing solution with cloth or sponge. **After using this compound, sanitized food surfaces must be thoroughly rinsed with potable water.**

PHENOL COEFFICIENTS:

Guaranteed	Minimum
S. aureus	30
S. typhosa	19

Rinse empty container thoroughly with water and discard.

LEM-O-DIS 19-30 is a cleaner, disinfectant, and sanitizer designed for general household and light industrial cleaning. For use on walls, floors and any inanimate hard surfaces. When used as directed,

CLEANER

DEODORIZER

LEMON ODOR

DISINFECTANT

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12,
 5% C18) dimethyl benzyl ammonium
 chlorides 1.6%
 n-Alkyl (68% C12, 32% C14) dimethyl
 ethylbenzyl ammonium chlorides 1.6%
 Sodium Carbonate 3.0%
 Essential Oils 0.2%

INERT INGREDIENTS 93.6%
 100.0%

DISINFECTANT

FUNGICIDE

LEM-O-DIS 19-30 will also deodorize those areas which generally are hard to keep fresh smelling, such as garbage pails, sink tops, and bathroom areas which are prone to odors caused by the growth of some micro-organisms. Effective against *Staphylococcus aureus* and *Salmonella choleraesuis* at 2 ounces per gallon. Also effective against pathogenic fungi, *Trichophyton interdigitale*, at this use dilution.

At 3 ounces per gallon, a hospital disinfecting claim against *Pseudomonas aeruginosa* can be made.

When used for sanitization of food processing and food contact items, limit the active quaternaries to 200 ppm. **After using this compound, sanitized food surfaces must be thoroughly rinsed with potable water.**

DANGER

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NET CONTENTS:

SEE RIGHT SIDE PANEL FOR ADDITIONAL
 PRECAUTIONARY AND FIRST AID STATEMENTS

DANGER

KEEP OUT OF REACH OF CHILDREN

MANUFACTURED FOR

A.B.C. COMPOUNDING CO., INC.,
 P. O. BOX 932
 ATLANTA, GEORGIA 30301

3862-53

3862-53

EPA REG NO 3862-53-
EPA EST NO. 3862-GA-1

LEM-O-DIS 19-30

USE DIRECTIONS

1. Add 2 ounces per gallon of this product to provide 500 ppm quaternary to the cleaning solution. This is enough for general disinfection, fungicidal activity against pathogenic fungi, and deodorizing.
For general deodorizing, LEM-O-DIS 19-30 is used at 200 ppm and applied to walls, sink tops, counter, and garbage pails.
2. For hospital disinfection per gallon, 1 ounce is required for disinfection against *Pseudomonas aeruginosa*.
3. Always follow proper cleaning steps recommended before using LEM-O-DIS 19-30.

Sanitization

When used for sanitization of previously cleaned food processing or food contact items, limit the level of active quaternaries to 200 ppm. 1/2 ounce per 1 gallon of water, and immerse articles in sanitizing solution or apply sanitizing solution with cloth or sponge. After using this compound, sanitized food surfaces must be thoroughly rinsed with potable water.

PHENOL COEFFICIENTS

Guaranteed	Minimum
S. aureus	30
S. typhosa	19

Rinse empty container thoroughly with water and discard.

LEM-O-DIS 19-30 is a cleaner, disinfectant, and sanitizer designed for general household and light industrial cleaning. For use on walls, floors, and any inanimate hard surfaces. When used as directed,

CLEANER

DISINFECTANT

DEODORIZER LEMON ODOR FUNGICIDE DISINFECTANT

ACTIVE INGREDIENTS

Active quaternary ammonium compound	1.4
Active quaternary ammonium compound	1.6
Active quaternary ammonium compound	3.0
Fragrance	0.2

INERT INGREDIENTS

93.6
100.0

SEE RIGHT SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY AND FIRST AID STATEMENTS

DANGER

KEEP OUT OF REACH OF CHILDREN

MANUFACTURED FOR

A.B.C. COMPOUNDING CO., INC.,
P. O. BOX 932
ATLANTA, GEORGIA 30301

LEM-O-DIS 19-30 will also deodorize those areas which generally are hard to keep fresh smelling, such as garbage pails, sink tops, and bath room areas which are prone to odors caused by the growth of some microorganisms. Effective against *Staphylococcus aureus* and *Salmonella choleraesuis* at 2 ounces per gallon. Also effective against pathogenic fungi, *Trichophyton interdigitale*, at this use dilution.

At 3 ounces per gallon, a hospital disinfecting claim against *Pseudomonas aeruginosa* can be made.

When used for sanitization of food processing and food contact items, limit the active quaternaries to 200 ppm. After using this compound, sanitized food surfaces must be thoroughly rinsed with potable water.

DANGER

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician if severe and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NET CONTENTS: