

BEST DOCUMENT AVAILABLE

ACCEPTED

M 1 21

30015

Active Thread :

Here: There: Only: .....

Keep eyes on road at all times

[illegible][illegible]

Do not reply in any way to this confidential  
letter without consulting the Bureau and the  
State of Washington.

Keep container closed until use. Product  
will freeze at temperatures below 0°C.

[illegible]

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total moisture content was determined by the method of AOAC (1990). The total dry matter content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).

BEST DOCUMENT AVAILABLE

... ..

BEST DOCUMENT AVAILABLE

longgrass, used at the same time as the annual weeds and grasses such as the  
bracken fern, Calluna, etc., and the  
restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.

Use the higher vegetation, etc., which is found in the  
bogs (such as the Calluna, etc.) and the

restiads, etc., which are found in the  
low-lying bogs. The plants are  
such as some of the ferns, etc., which  
are not the same as the annual weeds  
which are found in the bogs.