

U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
1200 Pennsylvania Ave, N.W.
Washington, D.C. 20460

NOTICE OF PESTICIDE:

Reregistration

x_Registration

EPA Reg. Number: 3862-133

Date of Issuance:

١٨٥

3 2001

Term of Issuance:

Conditional

Name of Pesticide Product;

Assault II (aerosol)

(under FIFRA, as a rended)

Name and Address of Registrant (include ZIP Code):

ABC Compounding PO Box 16247 Atlanta, GA 30321-0247

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is <u>conditionally</u> registered in accordance with FIFRA sec. 3(c)(7)(A), provided that you:

- 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of you product under FIFRA section 4.
- 2. Make the following labeling changes listed below before you release the product for shipment:
- a. Under your PRECAUTIONARY STATEMENTS section, please change the text to read "Harmful if absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse."
- b. Please move your STORAGE AND DISPOSAL section to the end of your label.

Signature of Approving Official:

Date

APR 3 2001

Commo T. La Dagga

George T. LaRocca Millia

EPA Form 8570-6

SPRM- | 4/3/01

- c. Under the Precautionary Statements section of your label, please move the animals paragraph to the directions for use on dogs.
- d. Either delete the statements "Repeat treatment as necessary", "Repeat as necessary", and "Repeat application as necessary" or change them to read "Repeat Every 4 Weeks", based upon the residual efficacy of the product.
- e. Please clarify which pests are controlled for four weeks based on the label claim "Controls Pests for Up to Four Weeks".
- f. Under the "Precautionary Statements" section, delete the statement beginning "If lice infestations should occur...etc.," since there are no EPA registered products labeled for this use on humans.
- 3. Please submit two (2) copies of your final printed label for before your release the product for shipment. Please refer to the A-79 enclosure for a further description of final printed labeling. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

If you have any questions regarding this action, please contact Dr. William Sproat of my team at (703) 308-8587.

Sincerely

George T. LaRocca

Product Manager (13)

Insecticide Branch

Registration Division (7505C)

Enclosure



- CONTAINS THE SYNTHETIC PYRETHROID PERMANONE®
- * CONTROLS NAMED RESIDENTIAL, INDUSTRIAL AND GARDEN INSECTS
- * FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS.
- * AN AQUEOUS STABLE MICROEMULSION
- CONTROLS PESTS FOR UP TO FOUR WEEKS
- KILLS LICE AND THEIR EGGS
- * FOR USE ON BEDDING, FURNITURE AND OTHER INANIMATE OBJECTS INFESTED WITH LICE.
- NOT FOR USE ON HUMANS
- * FOR USE ON LIVESTOCK AND THEIR PREMISES
- CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS

ACTIVE INGREDIENT:

100.00%

*(3-phenoxyphenyl)methyl (+/-) cis/trans 3-(2,2-dichloroethenyl)-2, 2-dimethykyclopropanecarboxylate (Cis/Trans ratio; Min. 35% (+/-) cis and max. 65% (+/-) trans.)

3862 - CGG

3862-GA-1

EPA REG. NO.

EPA EST. NO. 🐬

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF INHALED: Remove-affected person to fresh air. Apply artificial respiration if indicated.

IF IN EYES: Flush with plenty of water. Contact a physician if irritation persists.

IF ON SKIN: Wash affected areas immediately with soap and water. Get medical attention if irritation persists.

See Side Panel For Additional Precautionary Statements

ACCEPTED with COMMENTS to EPA Letter Dated

APR 3 2001

NET CONTENTS:

Under the Federal Inscaticide. Fungicide, and Rudenticida for an amended, for the production register 15 pages 13862 - 133



READ ENTIRE LABEL BEFORE EACH USE FOR PET USES: USE ONLY ON DOGS

PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals

CAUTION

Avoid spraying in eyes. Avoid breathing spray mist. Avoid contact with skin. Wash hands thoroughly after use. Vacate room after treatment and ventilate before reoccupying. Avoid contamination of feed and foodstuffs. Cover or remove fishbowls. This product is not for use on humans. If lice infestations should occur on humans, use an EPA registered product labeled for use on humans.

ANIMALS: Do not spray directly in/on eyes, mouth or genitalia of pets. Use only on dogs. Do not use on cats or kittens. Do not use on puppies less than twelve (12) weeks old. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated or nursing animals. Sensitivities may occur after using any pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Environmental Hazards

This product is toxic to fish. Do not apply to any body of water. Do not contaminate water when disposing of equipment washwaters.

Physical Or Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry area. Do not transport or store below 32"F. Disposal: Wrap container in several layers of newspaper and dispose of in trash. Can may be disposed of with other recyclables where local law permits. Do not incinerate or puncture.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not use in the edible product areas of food/feed processing plants, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as diring rooms but excluding areas where foods max be prepared or held.

in the home cover all food handling surfaces and cover or remove all food and cooking utensils, or wash thoroughly after treatment. Exposed food should be covered or removed.

NONFOOD AREAS: may be applied in nonfood/feed areas. Nonfood/feed establishment areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after canning or bottling).

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. Remove birds and cover fish aquariums before spraying.

HOUSEFLIES, FRUIT FLIES, MOSQUITOES, GNATS, WASPS, HORNETS, BEES, BLACK FLIES, SKIPPER FLIES, BLOW FLIES AND SMALL FLYING MOTHS: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which altract these insects. Keep nozzle in constant motion. Repeat treatment as necessary.

COCKROACHES, WATERBUGS, PALMETTO BUGS, SILVERFISH, FIREBRATS, CRICKETS, SCORPIONS, SPIDERS, MILLIPEDES, CENTIPEDES, SOWBUGS AND PILLBUGS: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible. Repeat treatment as necessary.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

INDIAN MEAL MOTHS, ANGOUMOIS GRAIN MOTHS: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application as necessary.

EXPOSED STAGES OF CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, RED FLOUR BEETLES, RICE WEEVILS, GRANARY WEEVILS, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

BEDBUGS: Spray mattresses, particularly around tufts and seams: Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Allow all sprayed articles to dry thoroughly before reuse. Repeat treatment as necessary.

TO CONTROL CLOTHES MOTHS AND CARPET BEETLES on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry

thoroughly before use.

IN ANIMAL QUARTERS AND MAK ROOMS: For rapid control of House Flies, Mosquitoes, Gnáts, Wesps, Hornets, Bees and Small-Flying Moths. Remove animals prior to application. Close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

TO TREAT LIVESTOCK: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

TO CONTROL BLOOD SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

FOR CONTROL OF BEDBUGS AND MITES IN POULTRY HOUSES, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

FLEAS AND TICKS: (ADULTS AND LARVAE): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

TO CONTROL FLEAS AND TICKS ON DOGS. Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application every two weeks if necessary.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use.

FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites. and Leafminers.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, Euonymus, fuschia, geraniums, crassulag grape vine, Kentia palm, lauret, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and rinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible, Repeat treatment as infestations occur.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions Disclaimer of Warranties before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application

All such risks shall be

assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: Seller warrants that this product corresponds to all claims and descriptions on the label thereof and is reasonably fit for use of any purpose stated on such label only when used in accordance with directions under normal use conditions. Seller makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.