



WARNING

Keep Away from Feed or Food Products

POISON

CAUTION—DO NOT DROP

IF LEAKING DON'T

Breathe Fumes
Touch Contents
Swallow

This is to certify that the contents of this package are properly described by name and are packed and marked and are in proper condition for transportation according to the regulations prescribed by the Interstate Commerce Commission.

FERGUSON FUMIGANTS, INC.
HAZELWOOD, MO.

Shipper's name required
hereon for ship-
ments by
EXPRESS

SAFETY PRACTICES

Care must be exercised by the user to prevent spillage of the fumigant on skin, clothing or shoes. If such spillage should occur, especially on clothing or shoes, such clothing or shoes should be immediately removed. Wash clothing and air shoes thoroughly before re-use. Repeated contact with the bare skin should be avoided and skin should be washed when wet with fumigant. In case of contact with eyes, flush with plenty of water for at least 15 minutes. **DO NOT WEAR GLOVES.** Do not use in container or applicator equipment with aluminum parts.

DO NOT REUSE EMPTY CONTAINER. Destroy by perforating or crushing and burying in a safe place. Use Eagle Claw opener only.

Keep away from heat. Keep container closed.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**WE
SELL
RESULTS**

RO

GELI

For Re
For the Ornament
CONTROLS Certain soil bor
and Stylet nematodes.



KEEP

WARNING: Poisonous Liquid
breathe vapor. Wear a full-face
Bureau of Mines for organic v
fact, immediately remove all
soap and water and flush eyes
diately. Do not reuse shoes or

Remove victim to fresh air im
ration if breathing has stoppe

ACTIVE I
Meth
Ethyl
INERT IN

The purchaser is licensed unde
fee. Additional Patent Pendin

Net Contents
USDA Reg. No. 3886-13

ROTOX

GELLED SOIL FUMIGANT

For Rotary Tillage Incorporation in Heavy Soils

For the Ornamental Nursery, Flower Growing and Turf Grass Industries.

CONTROLS Certain soil borne pests including certain weeds, weed seeds, Lesion, Spiral, Pin, and Stylet nematodes.



DANGER POISON



KEEP AWAY FROM CHILDREN

WARNING: Poisonous Liquid and Vapor! Contact with gelled liquid may produce burns. Do not breathe vapor. Wear a full-face gas mask with black canister meeting specifications of the U.S. Bureau of Mines for organic vapors. Do not get into eyes, on skin, or on clothing. In case of contact, immediately remove all contaminated clothing including shoes. Wash skin thoroughly with soap and water and flush eyes with water for at least 15 minutes. Get medical attention immediately. Do not reuse shoes or clothing until free of all contamination.

ANTIDOTE

Remove victim to fresh air immediately. Keep victim lying down and warm. Give artificial respiration if breathing has stopped. Call a physician immediately.


ACTIVE INGREDIENTS

Methyl Bromide

Ethylene Dibromide

INERT INGREDIENTS

One Gallon Weighs 5.3 Lbs.



The purchaser is licensed under U.S. Pat. No. 2,606,857 and the price paid includes the license fee. Additional Patent Pending.

Manufactured by

FERGUSON FUMIGANTS, INC.

95 Ford Lane, Hazelwood, Mo. 63042

Net Contents 5-

USDA Reg. No. 3886-12

Made in U.S.A.

Printed in U.S.A.

DIRECTIONS

ROTOX is to be used only as a preplant treatment.

Against Weeds, Nematodes — Use contrary to directions may result in injury to plants, unsatisfactory control or in other unintended consequences. Study entire label before use.

Soil preparation — The soil should be worked in advance if time is needed for cover crop, weed growth, or other plant refuse to decay before fumigation. Soil should have relatively high moisture content. Treat as soon as soil has dried out to tillable condition following rain or irrigation. This allows germination of weed seed and an activation of nematodes, thus increasing susceptibility to the fumigant.

BROADCAST FUMIGATION

FLORAL CROP BEDS, LANDSCAPE PLANT NURSERY BEDS, TURF GRASS SEED BEDS, SOIL ON WHICH TURF GRASS IS TO BE Laid use ROTOX at the rate of 20 gallons per acre in moist soil, at least 50°F. or above in temperature. Applicator equipment must release ROTOX from outlets spaced at 8 to 10 inch intervals at ground level near front or rotary tiller blades. Pull a roller behind the rotary tiller to compact soil and retard fumigant vapor loss. Allow 10 days between treatment and planting. A few types or species of bromine sensitive plants may require longer waiting periods such as Conifers, Delphinium, Holly, Multiflora Rose, Salvia and Snap Dragons. Carnations are extremely sensitive. Do not use this product for carnations. Adequate soil aeration is important. If there is a question, use a small plot to determine safety. ROTOX is not recommended for treatment of extremely light, sandy soil.

Rotary tiller should be set to penetrate a depth of 8 inches. Do not attempt fumigation at less than 6 inch depth. Consult your supplier for name and address of rotary tiller applicator attachment manufacturer.

To avoid re-infestation, do not use transplants, tools or crop residues that may be contaminated with soil borne pests.

READ ENTIRE LABEL BEFORE MAKING APPLICATION