

Nov 3, 1972
 UNDER THE FEDERAL INSECTICIDE,
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PESTS IN REGISTER
 ED NUMBER 3887-15
 TO ATTACHED COMMENTS

SECTROL DAIRY & MILL INSECTICIDE

DIRECTIONS

To kill the exposed stages of con-
 fused flour beetles, cadelltes, saw-
 toothed grain beetles, mediterranean
 flour moths, spider beetles, an-
 gousais grain moth and saw t-
 weevil, coffee bean weevil, rice
 weevil, lesser grain borer, foreign
 grain beetle, Mexican grain beetle,
 flour beetle, flat grain beetle, red
 horned grain beetle, raisin moth,
 tobacco moth, almond moth, cigar-
 ette beetle, drug store beetle,
 dried fruit beetle. Clean grain and
 flour bins, ciutes and elevators.
 Spray thoroughly and be sure to
 force insecticide into cracks and
 crevices in ceiling, wall and floor
 so as to penetrate the hiding
 places of the insects. Repeat
 application as required.

To control flies, gnats and mos-
 quitoes, in and at, creameries,
 dairies, food packing and food
 canning plants. Close doors and
 windows, create a dense fog in the
 room, leave the room closed at
 least 15 minutes, and sweep up
 dead insects.

Do not remain in treated areas.
 Ventilate the area after treatment
 is completed.

SECTROL DAIRY and MILL INSECTICIDE is specially manufactured for
 use in dairies, creameries, food packing plants, food processing
 plants, elevators, flour mills, feed mills and granaries, elevators
 and box cars.

SECTROL DAIRY and MILL INSECTICIDE is excellent for spraying in
 dairy barns and on cattle.

ACTIVE INGREDIENTS

Piperonyl butoxide, technical	100%
Pyrethrins	0.60%
Petroleum Distillate	0.12%
	99.28%

*Equivalent to 0.48% (butylcarbityl) (6-propylpiperonyl) ether and 0.12%
 related compounds.

CAUTION - KEEP OUT OF REACH OF CHILDREN.

- Avoid contamination of feed and foodstuffs.
- Do not spray while plant is in operation.
- FIRE HAZARD: Do not use or store near heat or open flame.

HARMFUL IF SWALLOWED

Manufactured by
Insect Control Sales and Service
Ephrata, Pa. 17522

NET CONTENTS: 5 GALS.

EPA REG. No. 3887-15

DIRECTIONS

Foods should be removed or cov-
 ered during treatment. All food
 processing surfaces should be
 covered during treatment or thor-
 oughly cleaned before using. When
 using the product in these areas
 apply only when the facility
 not in operation.

For cheese skippers, cheese mites
 and chocolate moths. Spray area
 thoroughly before storing. Force
 spray into cracks, crevices and
 hiding places of insects in floors,
 ceilings and walls. Repeat appli-
 cation as required.

For Cattle Spraying: To kill stable
 flies, horn flies, horse flies, deer
 flies and lice, both sucking and
 biting, spray directly on cattle,
 using a light application of not
 more than 1/2 or 2 ounces per
 adult animal. Avoid wetting hides
 of animals. Repeat application as
 required.

This product is toxic to fish. Keep
 out of lakes, streams, or ponds.
 Do not contaminate water by clean-
 ing of equipment, or disposal of
 wastes. Apply this product only
 as specified on this label.

Do not reuse empty container.
 Destroy it by perforating or crush-
 ing. Bury or discard in a safe
 place away from water supplies.