

A RICHMOND VETERINARY QUALITY PRODUCT

HEXRIN

DIRECTIONS

Mix 1½ ounces HEXRIN to one gallon water. Apply this HEXRIN emulsion to dog by sponge, spray, or dip. All surfaces should be thoroughly wetted. Allow emulsion to dry on dog.

Sold to Graduate Veterinarians Only
One U. S. Gallon

A concentrated dip or spray for the control of ticks, fleas, lice, and sarcoptic mange mites of dogs.

Active Ingredients:

Gamma Isomer of Benzene Hexachloride (From Lindane)	4.250%
*Technical Piperonyl Butoxide	3.950%
*(Equivalent to 3.159% of (butyl-carbityl) (6-propylpiperonyl) ether and 0.791% of related compounds.)	
Pyrethrins	0.395%
Pine Oil	10.000%
Polyoxyethylene Sorbitol mixed Ether Esters	14.810%
Petroleum Distillate	53.775%
Xylene	10.000%
	97.180%
Inert Ingredients	2.820%
	100.000%

See Right Panel for Additional Cautions

CAUTION—KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Use only as directed. Not for use on human beings or for household pets other than dogs. Avoid breathing spray mist. Avoid contact with skin or eyes. In case of contact with skin, wash with soap and water. Contact with skin can cause toxic symptoms. Avoid contamination of feed and foodstuffs. Toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Dispose of containers by flushing thoroughly with water and dispose of flush water where it will not contaminate feed, food or water source. Do not dip or spray dog more often than at 14 day intervals as protection against reinfestation.

USDA REG. No. 3890-2

For best results make up fresh emulsion and use immediately. Killing effect of Hexrin deteriorates rapidly after being in emulsion 12 hrs.

RICHMOND VETERINARY SUPPLY COMPANY
Distributors Richmond, Virginia 23225

