

ECHOLS RAT PELLETS

KILLS RATS AND MICE
CLEAN—EFFECTIVE
READY TO USE BAIT

Contains
WARFARIN!



CAUTION: Keep out of reach of children
NET WT. 1 LB. (See other cautions on back panel)

Sold by Athena Corporation
4838 Woodall, Dallas, Texas

ACCERTED

A detailed black and white illustration of a mouse, shown in profile facing left. It has a long tail, whiskers, and a small ear.

SEP 11 1972

**UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.**

FOR CONTROLLING RATS...
FOR CONTROLLING MICE...
IMPORTANT NOTICE...
CAL

ECHOLS RAT PELLETS

KILLS RATS AND MICE
CLEAN—EFFECTIVE
READY TO USE BAIT

Contains
WARFARIN!



CAUTION Keep out of reach of children
NET WT. 1 LB. (See other cautions on back panel)

Sold by **Athena Corporation**
Dallas, Texas

ACCEPTED
SEP 11 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.

3941-22

FOR CONTROLLING RATS Place one level spoonful of bait in each bait station. Bait stations should be placed in areas where rats are likely to feed. Refresh bait stations with fresh bait.

FOR CONTROLLING MICE Place one level spoonful quantity of bait in each bait station. Bait stations should be placed in areas where mice are likely to feed. Refresh bait stations with fresh bait. **IMPORTANT NOTICE:** More bait may be needed to control mice. Additional bait should be purchased as required.

CAUTION: Treated baits should be placed in locations not accessible to children, pets, wildlife or domestic animals. This material may be harmful if ingested. In such cases, intravenous and oral administration of Vitamin K₁ is recommended as in the case of hemorrhage caused by overdose of bait.
ACTIVE INGREDIENTS

Warfarin 3-(alpha-acetylbenzyl) 4-hydroxycoumarin
INERT INGREDIENTS
Total
Item No. E-16

AC
UNDE
FUNG

E.P.A. Reg. No. 3941-22

FOR CONTROLLING RATS: Place 4 to 6 ounces of bait in container where rats are known to feed or run. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Never let a bait station remain empty. Check daily. Replace stale, damp or moldy baits with fresh bait.

FOR CONTROLLING MICE: Place tablespoon quantities of bait in shallow containers where mice and signs of mice have been seen. Bait stations at intervals of 6 to 8 feet are recommended. Assure an interrupted supply of bait for a period of not less than 15 days, or until all signs of feeding have stopped.

IMPORTANT NOTICE: More bait may be needed to gain control than amount contained in this package. Additional bait should be purchased as required.

CAUTION:

Treated baits should be placed in locations not accessible to children, pets, wildlife, and domestic animals, or in tamper-proof bait boxes. Rinse empty container thoroughly with water and discard it. If swallowed by humans, pets, wildlife or domestic animals this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin.

ACTIVE INGREDIENTS

Warfarin 3-(alpha-acetylbenzyl)- 4-hydroxycoumarin 0.025%


INERT INGREDIENTS 99.975%

Total 100.00%

Item No. E-16

NET WT. 1 LB.

ACCEPTED



SEP 11 1972

**UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.**

3941-22

ACCEPTED

4

**UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 3941-22**