



# ECHOLS®

RAT and Mouse BAIT

# KILLS RATS


Contains

and Mice

# 4 Ready-to-Use BAIT PACKS

Net Weight: 1 lb  
4 (4 oz) Bait Packs

**CAUTION:** Keep out of reach of children. See back panel  
for additional cautions.



## DIRECTIONS FOR USE

Item No. E-1

**RATS:** For the control of Norway Rats and Roof Rats.

Open BAIT-PACKS and place 1 to 4 packages (4 to 16 ounces) in dry containers next to burrows, in runways, or where rats are accustomed to feed, or where signs of rats have been seen.

Bait should be placed in dark, out-of-the-way locations. Place along runways, behind boards, against wall.

Assure an uninterrupted supply of bait for a period of not less than fifteen days and continue baiting until all signs of feeding have stopped.

Replace stale, damp or moldy baits with fresh bait.

**MICE:** For the control of House Mice.

Open BAIT-PACKS and place tablespoonful amounts of bait ( $\frac{1}{4}$  to  $\frac{1}{2}$  ounce) in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of eight to twelve feet are recommended. Assure an uninterrupted supply of bait for a period of not less than fifteen days and continue baiting until all signs of feeding have stopped.

This rodenticide contains Warfarin, an anti-coagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice.

Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not re-use container. Destroy when empty.

### ACTIVE INGREDIENT

Warfarin 3 (alpha-acetonylbenzyl)-

4 hydroxycoumarin

INERT INGREDIENTS

TOTAL


0.025%

99.975%

100.000%

### CAUTION: Keep out of reach of children.

If swallowed by humans, domestic animals, wildlife or pets this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of Bishydroxycoumarin. In case of ingestion, call a physician.



ATHENA CORPORATION, 4838 Woodall, Dallas, Texas 75247

EPA Reg. No. 3941-27

EPA Est. No. 3941 TX 1

### New Coated Warfarin Bait!

Each particle of the Warfarin in this product has been coated with a material for the purpose of reducing detection of the Warfarin by the rodents at the time the bait is consumed. The coating may help assure greater acceptance of the bait.