

[4000-17]



PREMIUM FLY & ROACH SPRAY Grade AA Fly Spray

Active Ingredients: 100%
Petroleum hydrocarbons 99.38%, N-octyl Bicycloheptene Dicarboximide 0.34%, *Technical Piperonyl Butoxide 0.19%, Pyrethrins 0.09%.
Inert Ingredients: 0
*Equivalent to 0.152% (butylcarbityl) (6-propyl-piperonyl) ether and 0.038% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed.

Do not use, pour, spill or store near fire or open flame.

For additional precautions see also side panel.

Disributed By

APPROVED
JAN 7 1974
ESTABLISHED

MAR 6
UNDER
FUNGICIDE
FOR THE
ED GEN
TO APPL

KILLS ON CONTACT THE EXPOSED STAGES OF: Roaches, Waterbugs, Silverfish, Flies, Mosquitoes, Flying Moths, Bees, Hornets, Wasps, Gnats, Fruit Flies, Sand Flies, Weevils, Caddisflies, Grain Moths, Flour Beetles, Grain Mites, Stable Flies, Spiders. When used as directed, assures thorough destruction and control of insects infesting food processing, storage, handling or serving installations such as Flour Mills, Dairies, Meat Packing Plants, Grocery Stores, Restaurants, Barns, Grain Elevators, Warehouses. Can be used in all types of pump or atomizer dispensers adjusted to deliver a fine mist.

CAUTION

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation. Do not remain in treated areas. Ventilate the area after treatment is completed. Avoid inhalation of mist. Do not spray directly on plastics or asphalt tile. Do not reuse empty cans. Destroy by perforating or crushing and bury in a safe place away from water supplies. Do not reuse empty drum. Return to drum reconitioner or destroy by perforating or crushing and bury in a safe place away from water supplies.

DIRECTIONS

Roaches, Waterbugs, Silverfish: The infested areas should be sprayed thoroughly and repeatedly contacting as many insects as possible. Pay particular attention to hiding places, cracks, dark places, under sinks, behind pipes, etc.

Flies, Mosquitoes, Flying Moths, Bees, Hornets, Wasps, Gnats, Fruit Flies, Sand Flies: Close all windows and doors. Fill entire room with insecticide fog by directing spray toward ceiling. Leave room closed at least 10 minutes. Repeat as needed.

Weevils, Caddisflies, Grain Moths, Flour Beetles, Grain Mites: (Accessible stages) Apply in a wet, coarse spray to all infested surface areas, insect hiding places and runways, in flour mills, feed mills, warehouses. Exposed bags should be sprayed lightly. Repeat as necessary.

Control of Horn Flies and Stable Flies: For space spraying for control of flies when animals are on the premises, apply in a fine mist. In order to temporarily protect cows and horses from fly annoyance, spray animals from a distance of 6 to 8 feet using 1 to 2 ounces per animal daily or as necessary. Spray cows prior to milking taking care not to spray insecticide onto the udders. Wash udders with a sanitizing solution after spraying cow with insecticide prior to milking.

Fleas Infesting Premises: For quick and effective control, spray floors and cracks in the infested areas. Since fleas hide in upholstered furniture and carpets, these areas as well as wall cracks, baseboards and sleeping quarters of animals should be sprayed generously for direct contact with the pest.

Control of Red Mites and Blue Bugs in Poultry Houses: Remove all dirt and debris from infested area. Apply one quart of insecticide per 1,000 square feet of floor space. All surfaces and cracks should be wet thoroughly by spraying from a distance of 3 to 5 feet.

Spiders: Using a coarse wet spray, apply on or into spider webs covering area thoroughly. All visible Spiders should be wet with insecticide.

EPA REG. NO. 4000-19

NOTE: DO NOT SPRAY INSECTICIDE ONTO FOWLS

Mfg. By
SOUTHERN CHEMICAL PRODUCTS COMPANY
Macon, Georgia

REPTED

Mar. 6, 1974

4000-17