

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING**

**KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Causes skin irritation. Avoid contact with skin, eyes, and clothing.

**STATEMENT OF PRACTICAL TREATMENT**

**FIRST AID PROCEDURES:**

**EYES:** Flush with water for 15 minutes. Get medical attention.  
**SKIN:** Wash thoroughly with soap and water. Remove and wash contaminated clothing before reuse. **INGESTION:** Do not induce vomiting. Seek medical attention immediately.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Combustible mixture. Do not use, pour, spill or store near heat or open flame.

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

**CLEAN KILL**  
**SEASON LONG**  
**Non-Selective Weed Killer**

**ACTIVE INGREDIENTS:**

2,4-Dichlorophenoxyacetic acid, isooctyl ester	1.09%
Bromacil (5-bromo-3-secbutyl-1-methyluracil)	0.98%
<b>INERT INGREDIENTS:</b>	<b>97.93%</b>

\*2,4-Dichlorophenoic acid equivalent. TOTAL 100.00%  
0.72%

**WARNING**

**Harmful if swallowed.**  
**KEEP OUT OF REACH OF CHILDREN**  
See left panel for further precautions.

**1 U.S. GALLON/3.8 LITERS**

**STORAGE AND DISPOSAL**

- **STORAGE:** Keep product in original container. Do not put concentrate or dilute into food or drink containers. Do not contaminate feed, foodstuffs or drinking water. Do not store or transport near feed or food. Store at temperatures above 32°F.
- **PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Pesticide waste are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these waste cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
- **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**DIRECTIONS**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Kills on contact the exposed stages of: Roaches, waterbugs, silverfish, flies, mosquitoes, flying moths, bees, hornets, wasps, gnats, fruit flies, sand flies, weevils, cadellies, grain moths, flour beetles, grain mites, stable flies, spiders. When used as directed, assures thorough destruction and control of insects infesting food processing, storage, handling or serving installations such as flour mills, dairies, meat packing plants, grocery stores, restaurants, barns, grain elevators, warehouses. Can be used in all spray guns or atomizer dispensers adjusted to deliver a fine mist.

**For Food Preparation Areas:** Food should be removed or covered during treatment. Food contact surfaces and equipment must be washed with an effective cleaning compound followed by a potable water rinse

**DIRECTIONS (CONTINUED)**

**Roaches, waterbugs, silverfish:** The infested areas should be sprayed thoroughly and repeatedly contacting as many insects as possible. Pay particular attention to hiding places, cracks, dark places, under sinks behind pipelines, etc.

**Flies, mosquitoes, flying moths, bees, hornets, wasps, gnats, fruit flies and sand flies:** Close all windows and doors. Fill entire room with insecticide fog by directing spray toward ceiling. Leave room closed at least 15 minutes. Repeat as needed.

**Weevils, cadellies, grain moths, flour beetles, grain mites:** (Accessible stages) Apply in a wet, coarse spray to all infested surface areas, insect hiding places and runways, in flour mills, feed mills, warehouses. Exposed bags should be sprayed lightly. Repeat as necessary.

**Control of horn flies and stable flies:** For space spraying for control of flies when animals are on the premises, apply in a fine mist. In order to temporarily protect cows and horses from fly annoyance, spray animal from a distance of 8 to 8 feet using 1 to 2 ounces per animal daily or as necessary. Spray cows prior to milking taking care not to spray insecticide onto the udders. Wash udders with a sanitizing solution after spraying cow with insecticide prior to milking.

**Fleas infesting Premises:** For quick and effective control, spray floors and cracks in the infested areas. Since fleas hide in upholstered furniture and carpets, these areas as well as wall cracks, baseboards and sleeping quarters of animals should be sprayed generously for direct contact with the pest.

**Control of red mites and blue bugs in poultry houses:** Remove all dirt and debris from infested area. Apply one quart of insecticide per 1,000 square feet of floor space. All surfaces and cracks should be wet thoroughly by spraying from a distance of 3 to 5 feet.

**Spiders:** Using a coarse wet spray, apply on or into spider webs covering area thoroughly. All visible spiders should be wet with insecticide.

EPA Reg. No. 4000-19

EPA Est. 4000-GA-