



NET CONTENTS 1 U.S. GALLONS

# QUATERNARY AMMONIUM GERMICIDE CONCENTRATED

**ACTIVE INGREDIENTS:**

n-Alkyl (C <sub>14</sub> -50%, C <sub>12</sub> -40%, C <sub>16</sub> -10%)	
Dimethyl Benzyl Ammonium Chloride	10.00%
Ethanol	2.50%
Tetrasodium Salt of Ethylene Diamine	
Tetraacetic Acid	0.06%

**INERT INGREDIENTS:** 87.44%

Total 100.00%

## DANGER

**KEEP OUT OF REACH OF CHILDREN**

For additional precautions see also side panel.

Distributed By

## PHENOL COEFFICIENT

(A.O.A.C. Method, 20 Deg. C.)

*Staphylococcus aureus* - 40

*Salmonella typhosa* - 25

Non-inflammable, non-volatile, odorless

## DANGER

**KEEP OUT OF REACH OF CHILDREN:** Corrosive and skin irritation. Do not get in eyes, on skin or eyes and skin when handling. Harmful if swallowed or inhaled. Do not use for disinfection of food.

**FIRST AID:** In case of contact, immediately flush eye with water for at least 15 minutes. For eyes, call a physician. Wash contaminated clothing before reuse. If swallowed, drink large quantities of water. Call a physician.

**NOTE TO PHYSICIAN:** Probable mucosal damage. The use of gastric lavage. Measures against circulatory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.

**DIRECTIONS:** Do not mix with soap or synthetic detergents. Surfaces to be disinfected should first be washed and rinsed.

1. General disinfection in hospitals, public rooms, etc. Use 1 ounce of germicide per 4 gallons of water solution containing two ounces of germicide per gallon of water. Apply freely to surfaces.
2. Disinfection of barber shops and beauty parlors: Clean thoroughly, then soak combs, brushes and razors in a solution of 1 ounce of germicide per 1 gallon of water for 15 minutes.
3. Sanitization of glassware, silverware, dishes, etc.: Wash with a suitable detergent, rinse thoroughly, then immerse in a solution of 1 ounce of germicide in 4 gallons of water for 15 minutes. DO NOT USE A TOWEL.
4. Disinfection of rubber goods: Clean thoroughly with water, then immerse in a solution of 2 ounces of germicide per 4 gallons of water for 15 minutes.
5. Sanitization of equipment and utensils in food preparation: Clean thoroughly with water before using, remove or carefully protect food products. After using, equipment and utensils should be immersed in a solution of 1 ounce of germicide in 4 gallons of water for 15 minutes. Rinse with potable water before operations are resumed.

EPA REG. NO. 4000-51

Mfg. By  
SOUTHERN CHEMICAL PRODUCTS CO.  
Macon, Georgia

## PHENOL COEFFICIENTS

(A.O.A.C. Method, 20 Deg. C.):

Staphylococcus aureus - 40

Salmonella typhosa - 25

Non-inflammable, non-volatile, odorless

### DANGER

**KEEP OUT OF REACH OF CHILDREN:** Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard it.

**DIRECTIONS:** Do not mix with soap or synthetic detergents. Articles to be disinfected should first be washed and rinsed.

1. General disinfection in hospitals, public rooms and at home. Apply solution containing two ounces of germicide per 4 gallons of water freely to surfaces.
2. Disinfection of barber shops and beauty parlor equipment: Clean thoroughly, then soak combs, brushes and razors for 10 minutes in a solution of 1 ounce of germicide per 1 gallon of water.
3. Sanitization of glassware, silverware, dishes, cooking utensils: Clean with a suitable detergent, rinse thoroughly, then dip in a sanitizing rinse made with 1 ounce of germicide in 4 gallons of water, and air dry. **DO NOT USE A TOWEL.**
4. Disinfection of rubber goods: Clean thoroughly, then soak in solution diluted 2 ounces of germicide per 4 gallons of water for 10 minutes.
5. Sanitization of equipment and utensils in food processing plants: Before using, remove or carefully protect food products and packaging materials. After using, equipment and utensils must be thoroughly rinsed with potable water before operations are resumed.

EPA REG. NO. 4000-51

Mfg. By  
SOUTHERN CHEMICAL PRODUCTS COMPANY  
Macon, Georgia