

NET CONTENTS U.S. GALLON

DUAL QUAT DISINFECTANT BOWL CLEANER

ACTIVE INGREDIENTS:

| | |
|--|----------------|
| Octyl decyl dimethyl ammonium chloride | 1.250% |
| Dioctyl dimethyl ammonium chloride | 0.625% |
| Didecyl dimethyl ammonium chloride | 0.625% |
| Alkyl*amino betaine | 1.000% |
| Hydrogen chloride | 8.000% |
| INERT INGREDIENTS: | 88.500% |

TOTAL 100.000%

*Alkyl = C⁸-7%; C¹⁰-8%; C¹²-46%; C¹⁴-24%;
C¹⁶-10%; C¹⁸-5%

DANGER

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed.

See side panel for further precautions
and first aid

Manufactured by
SOUTHERN CHEMICAL PRODUCTS COMPANY, INC.
430 Lower Boundary St. • P.O. Box 205
Macon, Georgia 31297

ACCEPTED
05 SEP 1986
4000-91
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTRATION
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye and skin irritation. Avoid contact with skin, eyes, and clothing. Harmful if swallowed. Do not breathe vapor or fumes.

STATEMENT OF PRACTICAL TREATMENT

FIRST AID PROCEDURES

EYES: Flush with water for 15 minutes. Get prompt medical attention.

SKIN: Wipe off acid gently, immediately wash area with soap and water then cover with moist magnesia or baking soda. Call a physician if irritation persists.

INGESTION: Drink a teaspoonful or more of milk of magnesia or a cup of soapy water followed with milk or raw egg whites. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product, in concentrated form, is toxic to fish, birds, and other wildlife. Keep out of any body of water.

PHYSICAL OR CHEMICAL HAZARDS

Protect from freezing. Do not expose to temperatures below 32° F.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Flush toilet. Add 1 fluid oz. (2 tablespoons) directly to water in toilet bowl (¼ gallon). Brush to cover all bowl surfaces including under the rim. Flush toilet. For badly stained bowls, repeat above treatment using 2 to 4 ounces of bowl cleaner. For use in urinals, add 1 fluid oz. (2 tablespoons) to ¼ gallon of water. Apply by brush to all urinal surfaces. Flush.

At 1 fluid oz. per ¼ gallon water, this product is effective against *Pseudomonas aeruginosa*, *Salmonella choleraesuis* and *Staphylococcus aureus* (in toilet bowls and urinals). This performance has been confirmed by the AOAC Use-Dilution Test Procedure. The cationic detergent system of this bowl cleaner inhibits the acid corrosion of metal parts found around toilet bowls and urinals. This product is a disinfectant bathroom bowl cleaner designed for hospital use. It is especially formulated to kill many germs commonly found in hospital, industrial, and household environments. This product is designed for use in toilet bowls and urinals. This bowl cleaner contains an acid compatible detergent system that removes resistant stains and increases its disinfectant action. When used as directed, this product cleans, disinfects, and deodorizes in one easy step.

STORAGE AND DISPOSAL

STORAGE: Keep product in original container. Do not put concentrate or dilute into food or drink containers. Do not contaminate feed, foodstuffs or drinking water. Do not store or transport near feed or food. Store at temperatures above 32° F.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

EPA Reg. No. 4000-91

EPA Est. 4000-GA-1

| | | | |
|--|---|--|---|
| | | | miners will not be controlled by diazinon or similar type insecticides. |
| Peas (succulent only) | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites | 2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water | Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water | Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water | Broadcast to the soil just before planting. Immediately work into soil 4-8 inches. |
| NOTES: (1) Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom. | | | |
| Peppers | Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers | 2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest. |
| Potatoes | Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers | 2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |

| | | | |
|--|---|--|---|
| | | of 1 1/2 gals. of water | soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| Spinach | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Squash, Summer | Aphids, Thrips, Spider Mites | 2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. |
| Squash, Winter | Aphids, Thrips, Spider Mites | 2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest. |
| Sweet Corn | Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites | 2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water | Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water | Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water | Broadcast to the soil just before planting. Immediately work into soil 4-8 inches. |
| NOTE: To protect bees, do not apply to corn foliage during the pollen shed period. | | | |