

24-4-9

CONTAINS: UREA-FORM NITROGEN 2,4-D PLUS BANVEL D[†]

ACTIVE INGREDIENTS as herbicides:

Dimethylamine Salt of Dicamba (3,6 dichloro-o-anisic acid)*	28%
Dimethylamine Salt of related acids	05%
Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid**	6%
INERT INGREDIENTS relative to Herbicide Content	98.98%
TOTAL	100.00%

*Equivalent to .23% 3,6-dichloro-o-anisic acid
**Equivalent to .57% 2,4-dichlorophenoxyacetic acid
†BANVEL D TM of Velsicol Chemical Corporation

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE OTHER SIDE FOR ADDITIONAL CAUTIONS

**KILLS DANDELION- PLANTAIN-CHICKWEED
AND OTHER BROAD LEAF WEEDS**

FEEDS DESIRABLE GRASSES

GOLD TAG

**WEED
and
FEED**

Belgiano's

° GOLD TAG

WEED & FEED LAWN FOOD

24 - 4 - 9

8ep. 11-11/63

4007-2

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) - - - - -	24.00%
1.3% Ammoniacal Nitrogen derived from Urea-ammonia solution	
14.7% Water soluble nitrogen derived from Urea & Ureaformaldehyde	
8.0% Water insoluble nitrogen derived from Ureaformaldehyde	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅) - - - - -	4.00%
SOLUBLE POTASH (K ₂ O) - - - - -	9.00%
Potential acidity equivalent to 900 pounds calcium carbonate per ton	

BOLGIANO WEED & FEED contains lawn food of high Urea-Formaldehyde Nitrogen content, plus 2,4-D and Banvel D.

BOLGIANO WEED & FEED kills broadleaf weeds and

weed, plantain, buckhorn, burdock, mustard, nettle, jimson weed, lawn pennywort, pigweed, ground ivy and sow thistle.

CHICKWEED AND OTHER SIMILAR WEEDS CON-

TOTAL NITROGEN (N) - - - - - 24.00%
 1.3% Ammoniacal Nitrogen derived from Urea-ammonia solution
 14.7% Water soluble nitrogen derived from Urea & Ureaformaldehyde
 8.0% Water insoluble nitrogen derived from Ureaformaldehyde
 AVAILABLE PHOSPHORIC ACID (P₂O₅) - - - - - 4.00%
 SOLUBLE POTASH (K₂O) - - - - - 9.00%
 Potential acidity equivalent to 900 pounds calcium carbonate per ton

BOLGIANO WEED & FEED contains lawn food of high Urea-Formaldehyde Nitrogen content, plus 2,4-D and Banvel D.

BOLGIANO WEED & FEED kills broadleaf weeds and chickweed. It stimulates the growth of desirable turf-grasses.

BROADLEAF WEEDS CONTROLLED: Dandelion, rag-

weed, plantain, buckhorn, burdock, mustard, nettle, jimson weed, lawn pennywort, pigweed, ground ivy and sow thistle.

CHICKWEED AND OTHER SIMILAR WEEDS CONTROLLED: Common chickweed, mouse ear chickweed, little hop clover, bur clover, white clover, henbit, thyme-leaved speedwell, yarrow, knotweed, and spotted spurge.

DIRECTIONS

APPLY approximately 4 lbs. evenly over 1,000 sq. ft. (20' x 50') of lawn area. Contents of this bag will cover 5,000 sq. ft.

BROADLEAF WEEDS (such as those listed above): For good results application should be made when the temperature is above 60° F., and there is adequate moisture in the ground so that broadleaf weeds are in a growing state.

Chickweed may be controlled at the same time. Spring application is preferred but late summer or early fall application is also recommended to destroy weeds that have developed from seed during spring and summer.

CHICKWEED: Application may be made whenever the temperature is above freezing. Chickweed can be controlled at lower temperatures than broadleaf weeds. (A repeat application may be necessary.) When the temperature is close to or below freezing, immediate results will not be seen from the plant food in BOLGIANO Weed & Feed. However, the slow-release Urea-Formaldehyde (U-F) Nitrogen will be available when the growing conditions are such that grass can utilize it to advantage.

FOR BEST RESULTS APPLY WHEN GRASS IS MOIST.

MOWING: Grass should not be mowed for several days before and after application. Lawn seed may be sown 7 to 10 days after application of BOLGIANO Weed & Feed.

IMPORTANT: Avoid drift on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not apply within the drip line of trees or shrubs. Do not contaminate domestic or irrigation waters. Do not store in direct contact with fertilizers, seeds, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra, nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turfs.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact, flush with abundance of water. For eyes, get medical attention. Do not take internally. Keep out of the reach of children. Seller makes no warranty expressed or implied concerning the use of BOLGIANO Weed & Feed. Buyer assumes all risk of use or handling whether in accordance with directions or not. Do not store near feed or foodstuffs. Do not contaminate domestic or irrigation waters.

SPREADER SETTINGS

Modern	4½
Scott	5½
Sears	4½

F. W. BOLGIANO & CO.
WASHINGTON, D. C.

NET WT 20 POUNDS

Bolgianno's

GOLD TAG

WEED & FEED LAWN FOOD

24 - 4 - 9

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) ----- 24.00%

1.3% Ammoniacal Nitrogen derived from Urea-ammonia solution

14.7% Water soluble nitrogen derived from Urea & Ureaformaldehyde

8.0% Water insoluble nitrogen derived from Ureaformaldehyde

AVAILABLE PHOSPHORIC ACID (P₂O₅) ----- 4.00%

SOLUBLE POTASH (K₂O) ----- 9.00%

Potential acidity equivalent to 900 pounds calcium carbonate per ton

BOLGIANO WEED & FEED contains lawn food of high Urea-Formaldehyde Nitrogen content, plus 2,4 D and Banvel D.

BOLGIANO WEED & FEED kills broadleaf weeds and chickweed. It stimulates the growth of desirable turf-grasses.

BROADLEAF WEEDS CONTROLLED: Dandelion, rag-

weed, plantain, buckhorn, burdock, mustard, nettle, jimson weed, lawn pennywort, pigweed, ground ivy and sow thistle.

CHICKWEED AND OTHER SIMILAR WEEDS CONTROLLED: Common chickweed, mouse ear chickweed, little hop clover, bur clover, white clover, henbit, thyme-leaved speedwell, yarrow, knotweed, and spotted spurge.

DIRECTIONS

BOLGIANO

WEED & FEED LAWN FOOD

BEST DOCUMENT AVAILABLE

ACCEPTED
Sep. 3-1968
4067-2

24 - 4 - 9

GUARANTEED ANALYSIS

4067-3

TOTAL NITROGEN (N) - - - - -	24.00%
1.3% Ammoniacal Nitrogen derived from Urea-ammonia solution	
14.7% Water soluble nitrogen derived from Urea & Ureaformaldehyde	
8.0% Water insoluble nitrogen derived from Ureaformaldehyde	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅) - - - - -	4.00%
SOLUBLE POTASH (K ₂ O) - - - - -	9.00%
Potential acidity equivalent to 900 pounds calcium carbonate per ton	

BOLGIANO WEED & FEED contains lawn food of high Urea-Formaldehyde Nitrogen content, plus 2,4-D and Banvel D.

weed, plantain, buckhorn, burdock, mustard, nettle, jimson weed, lawn pennywort, pigweed, ground ivy and sow thistle.

BOLGIANO WEED & FEED kills broadleaf weeds and chickweed. It stimulates the growth of desirable turf-grasses.

CHICKWEED AND OTHER SIMILAR WEEDS CONTROLLED: Common chickweed, mouse ear chickweed, little hop clover, bur clover, white clover, henbit, thyme-leaved speedwell, yarrow, knotweed, and spotted spurge.

BROADLEAF WEEDS CONTROLLED: Dandelion, rag-

DIRECTIONS

10

APPLY approximately 4 lbs. evenly over 1,000 sq. ft. (20' x 50') of lawn area. Contents of this bag will cover 5,000 sq. ft.

MOWING: Grass should not be mowed for several days before and after application. Lawn seed may be sown 7 to 10 days after application of BOLGIANO Weed & Feed.

BROADLEAF WEEDS (such as those listed above): For good results application should be made when the temperature is above 60°F., and there is adequate moisture in the ground so that broadleaf weeds are in a growing state.

IMPORTANT: Avoid drift on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not apply within the drip line of trees or shrubs. Do not contaminate domestic or irrigation waters. Do not store in direct contact with fertilizers, seeds, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra, nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turfs.

Chickweed may be controlled at the same time. Spring application is preferred but late summer or early fall application is also recommended to destroy weeds that have developed from seed during spring and summer.

CHICKWEED: Application may be made whenever the temperature is above freezing. Chickweed can be controlled at lower temperatures than broadleaf weeds. (A repeat application may be necessary.) When the temperature is close to or below freezing, immediate results will not be seen from the plant food in BOLGIANO Weed & Feed. However, the slow-release Urea-Formaldehyde (U-F) Nitrogen will be available when the growing conditions are such that grass can utilize it to advantage.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact, flush with abundance of water. For eyes, get medical attention. Do not take internally. Keep out of the reach of children. Seller makes no warranty expressed or implied concerning the use of BOLGIANO Weed & Feed. Buyer assumes all risk of use or handling whether in accordance with directions or not. Do not store near feed or foodstuffs. Do not contaminate domestic or irrigation waters.

FOR BEST RESULTS APPLY WHEN GRASS IS MOIST.

SPREADER SETTINGS

BEST DOCUMENT AVAILABLE

Modern	4 1/2
Scott	5 1/2
Sears	4 1/2

WEED & FEED LAWN FOOD

24 - 4 - 9

8ep. 1968

4067-3

4067-3

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) - - - - - 24.00%
 1.3% Ammoniacal Nitrogen derived from Urea-ammonia solution
 14.7% Water soluble nitrogen derived from Urea & Ureaformaldehyde
 8.0% Water insoluble nitrogen derived from Ureaformaldehyde
 AVAILABLE PHOSPHORIC ACID (P₂O₅) - - - - - 4.00%
 SOLUBLE POTASH (K₂O) - - - - - 9.00%
 Potential acidity equivalent to 900 pounds calcium carbonate per ton

BOLGIANO WEED & FEED contains lawn food of high Urea-Formaldehyde Nitrogen content, plus 2,4-D and Banvel D.

BOLGIANO WEED & FEED kills broadleaf weeds and chickweed. It stimulates the growth of desirable turf-grasses.

BROADLEAF WEEDS CONTROLLED: Dandelion, rag-

weed, plantain, buckhorn, burdock, mustard, nettle, jimson weed, lawn pennywort, pigweed, ground ivy and sow thistle.

CHICKWEED AND OTHER SIMILAR WEEDS CONTROLLED: Common chickweed, mouse ear chickweed, little hop clover, bur clover, white clover, henbit, thyme-leaved speedwell, yarrow, knotweed, and spotted spurge.

DIRECTIONS

APPLY approximately 4 lbs. evenly over 1,000 sq. ft. (20' x 50') of lawn area. Contents of this bag will cover 5,000 sq. ft.

BROADLEAF WEEDS (such as those listed above): For good results application should be made when the temperature is above 60°F., and there is adequate moisture in the ground so that broadleaf weeds are in a growing state.

Chickweed may be controlled at the same time. Spring application is preferred but late summer or early fall application is also recommended to destroy weeds that have developed from seed during spring and summer.

CHICKWEED: Application may be made whenever the temperature is above freezing. Chickweed can be controlled at lower temperatures than broadleaf weeds. (A repeat application may be necessary.) When the temperature is close to or below freezing, immediate results will not be seen from the plant food in BOLGIANO Weed & Feed. However, the slow-release Urea-Formaldehyde (U-F) Nitrogen will be available when the growing conditions are such that grass can utilize it to advantage.

FOR BEST RESULTS APPLY WHEN GRASS IS MOIST.

MOWING: Grass should not be mowed for several days before and after application. Lawn seed may be sown 7 to 10 days after application of BOLGIANO Weed & Feed.

IMPORTANT: Avoid drift on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not apply within the drip line of trees or shrubs. Do not contaminate domestic or irrigation waters. Do not store in direct contact with fertilizers, seeds, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra, nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turfs.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact, flush with abundance of water. For eyes, get medical attention. Do not take internally. Keep out of the reach of children. Seller makes no warranty expressed or implied concerning the use of BOLGIANO Weed & Feed. Buyer assumes all risk of use or handling whether in accordance with directions or not. Do not store near feed or foodstuffs. Do not contaminate domestic or irrigation waters.

SPREADER SETTINGS

Modern 4½
 Scott 5½
 Sears 4½

BEST DOCUMENT AVAILABLE



CONTAINS: UREA-FORM NITROGEN

2,4-D PLUS BANVEL D¹

ACTIVE INGREDIENTS as herbicides:

Dimethylamine Salt of Dichloro-3,4-dinitrobenzoic acid

Dimethylamine Salt of related acids

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid

INERT INGREDIENTS relative to Herbicide Content

TOTAL 100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE OTHER SIDE FOR ADDITIONAL CAUTIONS

¹Equivalent to .23% 3,6-dichloro-2-nitrobenzoic acid
²Equivalent to .57% 2,4-dichlorophenoxyacetic acid
BANVEL D TM of Valent Chemical Corporation

BEST DOCUMENT AVAILABLE

KILLS

DANDELION-PLANTAIN-CHICKWEED AND OTHER BROAD LEAF WEEDS

FEEDS

DESIRABLE GRASSES

GOLD TAG

WEED and FEED

USDA REG. NO. 4067-3