


AUTHORIZED FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS

ACTIVE-INGREDIENTS:
Pyrethrins
Technical Piperonyl butoxide*
Petroleum Solvent
*Equivalent to 1.5% (butylcarbityl)
(6-propylpiperonyl) ether and 0.4%
related commounds.

related compounds.
INERT INCREDIENTS:

Net Contents 8 az. Avdp. EPA Reg. No. 4077 AC EPA Est. No. 4077 PA 1



Manufactured By:

ORB INDUSTRIES, INC. #2 Race St., P. O. Box 1067 Upland, PA 19015 ...

PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS TION

PHYSICAL OR CHEMICAL HAZARL flammable. Contents under pressure. Do not use near fire, sparks, or flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

CAUTION Harmful if swallowed or absorbed through skin. Avoid breathing vapors or sprey mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Reep out of reach of children. Ventilate after treatment is completed.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. In the home, all food processing surfaces and utensits should be covered during treatment, or thoroughly washed before use.

Remove pets, birds, and cover fish aquariums before spraying.

Storage and Disposal

Storage. Store in a cool dry area away from heat or open flame. Disposal. Do not reuse empty container. Wrap container and put in trash collection.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in home, institutions, foud processing plants, réstaurants, kitchens, dairies, closed barns, warehouses, box cars and trucks. Authorized for use in federally inspected meat and poultry plants.

and poultry plants.

ROACHES: Open drawers, cabinets and similar places and direct mist into such places as well as into cracks, crevices and other potential harbourages in order to activate insects, using a dosage of 30 seconds per 100 square feet of floor space. Spray insects directly when possible. Use conveneint ORB 4114 X Extension Nozzle for crack and cravice treatment. Follow immediately by applying as a space insecticide. Fill the room with spray mist, spraying 8 to 5 seconds per 1000 cubic feet of space. Sweep up fallen insects. Keep room closed for 30 minutes after treatment. Repeat treatment weekly until infestation is under control and then every two weeks or as needed.

