

PROPOSED REVISED LABEL

50 LBS. NET

ARBIE RUMINANT SCOOTER FORMULA R-B-N MINERAL FEED

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle

ACTIVE INGREDIENT	BY WEIGHT
chloro-1- (2, 4, 5-trichlorophenyl) .inyl dimethyl phosphate.	1.24%
INERT INGREDIENTS	98.76%**
Total 100.00%	

*Each pound contains 5.6 gms RABON.
**Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Calcium (Ca) not less than	16.50%
Calcium (Ca) not more than	18.00%
Phosphorus (P), not less than	5.50%
Salt (NaCl) not less than	9.50%
Salt (NaCl) not more than	10.50%
Iodine (I), not less than	0.056%
Vitamin A (as retinol), U.S.P. units per pound	

Dicalcium Phosphate, Ground Limestone, Sodium Bicarbonate, Salt, Wheat Middlings, Dried Distillers' Fermented Corn Extractives, Condensed Fish Solubles, Carbonate, Sulfur, Oil of Anise, Potassium Iodide, Vitamin A Acetate (with improved stability), Vitamin E Supplement, Vitamin D-3 Supplement (with improved stability), Menaquinone Dimethylpyrimidinol Bisulfite, Copper Oxide, Cobalt Carbonate, Manganous Oxide, Zinc Oxide, Cane Molasses

CAUTION - Keep Out Of Reach Of Children
CAUTION - Hazardous To Humans

Precautionary Statements
Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

General Classification

It is a violation of Federal law to use this product in a manner inconsistent with its labelling.

Feed Arbie Ruminant Scooter Formula R-B-N Mineral Feed at the rate of one ounce per 500 lbs. of body weight per head per day by mixing it into the ground grain or complete daily ration, or by feeding free choice (do not feed free choice while cattle are receiving Arbie Ruminant Scooter Formula R-B-N Mineral Feed or other source of Rabon mixed in the ground grain or complete daily ration).

Feeding Free Choice - Cattle should consume an average of 6 oz. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON® program but OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head, and place near watering and loafing areas. Put out no more than one month's supply at a time, and protect from rain.

If consumption of Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON® is above 6 oz. per 100 lbs. of body weight per month, reduce the number of feeding locations or rations. If consumption is below 6 oz. per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON® early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Arbie Ruminant Scooter Formula R-B-N Mineral Feed may be fed up to slaughter and to lactating dairy cows.

Arbie Ruminant Scooter Formula R-B-N Mineral Feed prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not as effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.


In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Arbie Ruminant Scooter Formula R-B-N Mineral Feed should be used in conjunction with other good management practices.

Storage and Disposal

Do not contaminate water or food by storage or disposal. Dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place.

Manufactured By
ARBIE MINERAL FEED CO., INC.
404 S. Center Street
EPA Reg No. 4089-5 EPA EST No. 04405 1A-A
Made in USA



71

72

ARBIE RUMINANT SCOOTER FORMULA R-B-N MINERAL FEED

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,6-trichlorophenyl) vinyl dimethyl phosphate.	1.24%
INERT INGREDIENTS	98.76%**
Total 100.00%	

*Each pound contains 5.6 gms RABON
**Refers only to ingredients which are not larvicides.

GUARANTEED ANALYSIS

Calcium (Ca), not less than	16.50%
Calcium (Ca), not more than	18.00%
Phosphorous (P), not less than	5.50%
Salt (NaCl), not less than	9.50%
Salt (NaCl), not more than	10.50%
Iodine (I), not less than	0.056%
Vitamin A	100,000 U.S.P. units per pound

FEED INGREDIENTS

Dicalcium Phosphate, Active Dry Yeast, Ground Limestone, Sodium Sulfate, Sodium Bicarbonate, Salt, Wheat Middlings, Dried Condensed Fermented Corn Extractives, Condensed Fish Solubles, Iron Carbonate, Sulfur, Oil of Anise, Potassium Iodide, Vitamin A Acetate (with improved stability), Vitamin E Supplement, Vitamin D-3 Supplement (with improved stability), Menadione Dimethylpyrimidinol Bisulfite, Copper Oxide, Cobalt Carbonate, Manganous Oxide, Zinc Oxide, Cane Molasses.

CAUTION — Keep Out of Reach of Children
CAUTION — Hazardous To Humans

Precautionary Statements

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

General Classification

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed Arbie Ruminant Scooter Formula R-B-N Mineral Feed at the rate of one ounce per 500 lbs. of body weight per head per day by mixing it into the ground grain or complete daily ration, or by feeding free choice (do not feed free choice while cattle are receiving Arbie Ruminant Scooter Formula R-B-N Mineral Feed or other source of Rabon mixed in the ground grain or complete daily ration).

Feeding Free Choice — Cattle should consume an average of 8 oz. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg RABON* Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON* program but OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head, and place near watering and loafing areas. Put out no more than one month's supply at a time, and protect from rain.

If consumption of Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON* is above 8 oz. per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 8 oz. per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON* early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Arbie Ruminant Scooter Formula R-B-N Mineral Feed may be fed up to slaughter and to lactating dairy cows.

Arbie Ruminant Scooter Formula R-B-N Mineral Feed prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not as effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter on the premises.

In order to achieve optimum fly control, Arbie Ruminant Scooter Formula R-B-N Mineral Feed should be used in conjunction with other good management practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or in a way approved for pesticide control. Do not bury.

Manufactured by

ARBIE MINERAL FEED CO., INC.

404 S. Center Street
EPA Reg. No. 4089-5 EST No. 04089-1A-A
Made in USA

10/73

PREPARED BY THE D. L. L. CO.

50 LBS. NET

ARBIE RUMINANT SCOOTER FORMULA R-B-N MINERAL FEED

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle

ACTIVE INGREDIENT	BY WEIGHT
1-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate.	1.24%
INERT INGREDIENTS	98.76%**
Total 100.00%	

*Each pound contains 5.6 gms RABON
**Refers only to ingredients which are not larvicidal

GUARANTEED ANALYSIS	
Calcium (Ca), not less than	16.50%
Calcium (Ca), not more than	18.00%
Phosphorous (P), not less than	5.50%
Salt (NaCl), not less than	19.50%
Salt (NaCl), not more than	10.50%
Iodine (I), not less than	0.056%
Vitamin A as retinol, 100,000 U.S.P. units per pound	

INGREDIENTS: Dicalcium Phosphate, Active Dry Yeast, Ground Limestone, Sodium Sulfate, Sodium Bicarbonate, Salt, Wheat Middlings, Dried Distillers' Solubles, Fermented Corn Extractives, Condensed Fish Solubles, Iron Carbonate, Sulfur, Oil of Anise, Potassium Iodide, Vitamin A Acetate (with improved stability), Vitamin E Supplement, Vitamin D-3 Supplement (with improved stability), Menadione Dimethylpyrimidinol Bisulfite, Copper Oxide, Cobalt Carbonate, Manganese Oxide, Zinc Oxide, Cane Molasses.

CAUTION - Keep Out Of Reach Of Children

CAUTION - Hazardous To Humans
Precautionary Statements

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

General Classification

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed Arbie Ruminant Scooter Formula R-B-N Mineral at the rate of one ounce per 500 lbs. of body weight per day by mixing it into the ground grain or complete daily ration or by feeding free choice (do not feed free choice while cattle are receiving Arbie Ruminant Scooter Formula R-B-N Mineral Feed or other source of Rabon mixed in the ground grain or complete daily ration)

Feeding Free Choice — Cattle should consume an average of 6 oz. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg. HABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON® program but OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head and place near watering and loafing areas. Put out no more than one month's supply at a time and protect from rain.

If consumption of Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON® is above 6 oz. per 100 lbs. of body weight per month, reduce the number of feeding locations or rate. If consumption is below 6 oz. per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Arbie Ruminant Scooter Formula R-B-N Mineral Feed with RABON® early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall.

Arbie Ruminant Scooter Formula R-B-N Mineral Feed may be fed up to slaughter and to lactating dairy cows.

Arbie Ruminant Scooter Formula R-B-N Mineral Feed prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not as effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control horn, house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Arbie Ruminant Scooter Formula R-B-N Mineral Feed should be used in conjunction with other good management practices.

Storage and Disposal
Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in a incinerator or landfill approved for pesticide containers or in a safe place.
Manufactured By
ARBIE MINERAL FEED CO., INC.
404 S. Center Street, Marshalltown, Iowa 50158
EPA Reg. No. 4089-5 EPA EST. No. 04085-1A
Made in USA 5 86