



Dec 10, 1960

4090-131
EPA REG. NO.

POST-EMERGENCE WEED KILLER EMULSIFIABLE LIQUID

(Contains 1 pound active ingredient per gallon)

ACTIVE INGREDIENTS:	12.8%
Cyprazine [2-chloro-4-(cyclopropylamino)-6-(isopropylamino)-s-triazine]	12.2% (min.)
Related compounds	0.6% (max.)
INERT INGREDIENTS:	87.2%
	100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Use Care in Handling. Avoid Contact with Eyes, Skin, and Clothing. Avoid Inhalation of Vapor. In case of Contact with Skin or Clothing, Wash with Soap and Water. If Eyes are Affected, Wash with Water.

NET CONTENTS: 5 GALLONS

EPA Registration No. 4090-131-AA

GULF OIL CHEMICALS COMPANY
Division of Gulf Oil Corporation, Houston, Texas 77002 U.S.A.

STIR WELL BEFORE USING:

General Information

**For Weed Control in Field, Sweet, and
Silage Corn and Popcorn**

OUTFOX is a selective herbicide recommended for post-emergence use on corn.

OUTFOX controls the following seedling weeds:

ANNUAL BROADLEAF WEEDS

Buffalo bur	Lambsquarters	Puncturevine
Burdock	Annual morningglory	Ragweed
Caryopside	Wild mustard	Smartweed
Cocklebur	Black nightshade	Sunflower
Flower-of-an-hour	Pigweed	Velvetleaf
Jimsonweed	Prickly lettuce	Tall waterhemp
Kochia	Prickly sida	Wild buckwheat

ANNUAL GRASSES

Barnyardgrass	Green foxtail
Downy brome	Yellow foxtail
Crabgrass	Shattercane
Bristly foxtail	Wild oats
Giant foxtail	Witchgrass

To avoid possible injury, do not allow spray drift of OUTFOX near ornamental trees, shrubs, or plants.

Slight temporary leaf burn or stunting may occur if corn is treated when wet or when under stress from cold weather or poor fertility.

To treat weeds or grass which have escaped a previous herbicide application, OUTFOX may be applied using either drop nozzles or a spray directed to avoid contact with the corn plant. Injury may occur if OUTFOX is applied directly to the foliage of corn previously treated with a pre-emergence or post-emergence herbicide.

Do not cut for feed or silage within 30 days after application of OUTFOX.

Following harvest of a treated crop, plow and thoroughly till the soil in fall or spring to avoid possible injury to spring-seeded crops other than corn.

Do not plant treated areas to any crop except corn until the following year. In California delay planting any crop except corn in OUTFOX-treated areas for at least eighteen months following treatment.

Do not plant sugar beets on land treated the previous year with OUTFOX.

Avoid contamination of food and feed.

Adding insecticides, herbicides, liquid fertilizers or other materials to OUTFOX is not recommended because of possible compatibility problems which could cause crop injury.

IMPORTANT NOTICE—Conditions of Sale: GULF warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. GULF neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation express or implied, concerning this material. Critical and unforeseeable factors beyond GULF's control prevent it from eliminating all risks in connection with the use of this material. Such risks include, but are not limited to, damage to plants and crops to which the material is applied. Such risks occur even though the material is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by GULF in the preceding paragraph) resulting from handling, storage, and use of the material.

Directions For Use

To determine the height of a grassy weed, extend the plant vertically to its full length and measure the distance between the tip of the longest leaf and the soil surface.

Optimum results are obtained by treating rapidly growing plants. Older weeds not growing rapidly, including those which are less than 2 inches tall (for California 1½ in.) may not be satisfactorily controlled.

Do not apply after corn is 10 inches high.

Before use thoroughly clean all spray equipment, nozzles, screens, and filters. Check the pump and pressure gauge to ensure that they are working properly.

For Areas Other Than California

For broadcast treatment apply OUTFOX at a spray volume of 10-30 gal/A and a pressure of at least 30 pounds at the following rate:

Grassy Weed Height, (inches)	OUTFOX (Quarts/A)
less than 2	3

For band application reduce the rate proportionately.

For California Only

For broadcast treatment apply OUTFOX at a spray volume of 10-30 gal/A and a pressure of at least 30 pounds at the following rate:

Grassy Weed Height, (inches)	OUTFOX (Quarts/A)
less than 1½	4

For band application reduce the rate proportionately.

Treat only rapidly growing weeds and grasses when the top soil moisture is sufficient to sustain rapid growth for several days. Spraying under adverse conditions such as heat, drought and low humidity conditions, etc. will produce less than acceptable weed control. Do not plant treated areas to any crop except corn until at least 18 months following OUTFOX treatment.

IMPORTANT

**STIR OUTFOX WELL BEFORE USING. NEVER POUR OUTFOX
INTO AN EMPTY SPRAY TANK. MAINTAIN GOOD AGITATION
IN SPRAY TANK WHILE ADDING OUTFOX.**


Partially fill the spray tank with about one-half the amount of water required, and begin good agitation. Normal bypass agitation is usually satisfactory. Return lines to tank should discharge below the liquid level. Stir the OUTFOX well and pour it slowly into the tank, maintaining good agitation during mixing and, subsequently, during filling, and spraying. Rinse the can thoroughly with clear water and complete the filling of the tank.

For optimum weed control after treatment, a timely shallow cultivation is recommended.

**DO NOT RE-USE EMPTY CONTAINER.
WASH THOROUGHLY WITH WATER AND DETERGENT.
DISCARD IN A SAFE PLACE.**

Manufactured and sold under U.S. Patents 3,451,802 and 3,503,971

C1013C-L-LC



MADE IN U.S.A.

EPA Est. No. 4090KS01

LITHO IN U.S.A.