

EL BLANCO
BRAND

TERRACLOR* THIMET 6.5 - 6.5

SOIL FUNGICIDE - SYSTEMIC INSECTICIDE GRANULAR

REG TRADEMARK OF AMERICAN CYANAMID COMPANY

*TERRACLOR IS A REGISTERED TRADEMARK OF
OLIN MATHIESON CHEMICAL CORPORATION

ACCEPTED
OCT 23 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FERTILIZER ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 4110-42 SUBJECT
TO ATTACHED COMMENTS.

10656-29

Active Ingredients:
Fenitrothion, O,O-diethyl S-(methylthio)methyl phosphorothioate, 6.50%
Thiometon-methyl, 6.50%
Inert Ingredients, 87.0%

DIRECTIONS FOR USE ON COTTON

For the control of early season insects (aphids, black cutworms, leafhoppers, leafminers, thrips, whiteflies and mites) and seedling disease complex-Rhizoctonia, Pythium and Fusarium on cotton, use 15 pounds per acre (40-inch row spacing) or 18 ozs. per 1,000 ft. of row (for any row spacing). Distribute evenly in the furrow at planting time.

USA Reg no. 4110-

DANGER

IF SWALLOWED, RAPIDLY ABSORBED THROUGH SKIN.
SYMPTOMS PROGRESSIVELY INCREASE SUSCEPTIBILITY TO POISONING.



**DANGER
POISON**



**KEEP AWAY
FROM
CHILDREN**

DO NOT EAT FOOD OR

DRINK BEVERAGES
OR USE TOBACCO
WHILE USING THIS
PRODUCT OR
FOR SEVERAL
DAYS THEREAFTER.
WASH YOUR HANDS
THOROUGHLY WITH
SOAP AND WATER
AFTER EACH USE.

DO NOT BLEACH
CLOTHING OR
OTHER FABRICS
WHICH HAVE
CONTACT WITH
THIS LABEL.

IF ON SKIN OR CLOTHING,

REMOVE IMMEDIATELY
BY WASHING WITH
SOAP AND WATER.

IF IN EYES,
WASH IMMEDIATELY
WITH COLD WATER
FOR SEVERAL
MINUTES.

IF INGESTED,
DO NOT INDUCE
VOMITING. CALL
YOUR PHYSICIAN
IMMEDIATELY.

IF SWALLOWED,
DRINK MILK OR
Egg whites.

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a supply of 1/100 grain tablets for emergency use. Never take atropine unless symptoms of poisoning have occurred.

FIRST AID: If symptoms or signs of poisoning (See Physicians Note) include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1/100 grain) at once. Consult a physician at once in all cases of suspected poisoning.

Remove patient immediately from the area. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of the pesticide. If swallowed, induce vomiting by giving warm salty or soapy water. Repeat until vomit fluid is clear. Have patient lie down and keep quiet. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

PHYSICIANS NOTE: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Give atropine intramuscularly or intravenously, 2 to 4 milligrams (3 to 6 tablets of 1/100 grain each) at once and every hour as required until pupils dilate. There may be 2 milligrams, or more may be required during the first 24 hours. Never give barbiturates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, increase susceptibility to very small doses of any cholinesterase inhibitor. Repeat atropine tests until cholinesterase regeneration has taken place as indicated by blood tests.


Buyer hereby certifies that the material contained herein is true and correct and is reasonably true to the use therein described. The manufacturer's warranty shall be limited to direct damages, and shall not include consequential damages such as loss of profits or values.

SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY. Buyer's use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by MANUFACTURER and SELLER.

Manufactured by

WHITE CHEMICAL COMPANY

1310 W. WATKINS ROAD • PHOENIX, ARIZONA 85007



Cameraman's Note
Poor Copy