

ACCEPTED
JUN 5 1972
UNDER THE FEDERAL DIRECTORATE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO.

CONTENTS 50 LBS. NET

RED KAY 5% SEVIN* DUST

4139-26

ACTIVE INGREDIENT : Carbaryl (1-naphthyl N-Methylcarbamate) 5%
INERT INGREDIENTS 95%
*Trade-Mark of UCC for 1-Naphthyl N-Methylcarbamate

CAUTION: KEEP OUT OF REACH OF CHILDREN

DIRECTIONS FOR USE

PEANUTS—For Control of Thrips and Leafhopper.

Thrips. Apply at the rate of 20 lbs. per acre using a regular duster with nozzles directly over the Peanut plants. One dusting application usually gives satisfactory control. Apply about the first or second week in June, or as soon as injury is first observed. May be applied up to and including the day of harvest.

Leafhopper. Apply dust at the rate of 20 lbs. per acre, using regular duster with nozzles directly over the Peanut plants. Two applications, approximately two weeks apart, usually needed. The first application should be made about the middle of July, or as soon as injury is first observed, and a second application should be made about the first of August. Additional applications may be made until the day of harvest.

TOBACCO—For the Control of Hornworms, Budworms, Flea Beetles, June Beetles and Japanese Beetles.

Applications should begin at the first sign of insect or their damage. Apply formulation at the rate of 25 lbs. per acre. Repeat application as needed. Do Not apply within three days of priming or cutting. Avoid excessive applications. If late season applications are necessary, treat immediately after priming rather than before priming.

BEE CAUTION: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CAUTION: Harmful if swallowed or inhaled. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands before eating. Wear regular long sleeved work clothing. Take shower or bath after work. Change to clean clothing daily. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and water receptacles. Do not reuse container. Destroy when empty.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible, Cholinesterase, inhibitor. Atropine is antidotal.

NOTICE: Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

EPA Reg. No. 4139-26

Meherrin Agricultural & Chemical Company
SEVERN, NORTH CAROLINA