

USE DIRECTIONS

Household

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and shower stalls. Apply solution with mop or cloth. Sanitize with 1 ounce Ark San to 4 gallons of water. Disinfect with 3 ounces Ark San to 5 gallons of water. Kills staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Hospitals

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use Ark San at 3½ ounces per 5 gallons of water. Apply with cloth, mop or sponge. At this level Ark San will be effective against Pseudomonas aeruginosa PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Food Processing Equipment

For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce Ark San per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, NO POTABLE WATER RINSE is required.

Dairies

To sanitize dairy equipment such as tanks, lines, pails and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce Ark San to 4 gallons of water (200 ppm). At this level, NO POTABLE WATER RINSE is required. Follow recommendations of local Health Board.

Restaurant and Bar Rinse

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce Ark San to 4 gallons of water. NO TERMINAL POTABLE WATER RINSE REQUIRED.

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per 5 gallons of water is effective against influenza A, Herpes Simplex, Adenovirus type 2, and Vaccinia Viruses.

ARK SAN

Disinfectant, Deodorizer & Sanitizer

Active Ingredients:

| | |
|---|------------|
| n-Alkyl (60% C ₁₁ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₃) dimethyl benzyl ammonium chlorides | 5% |
| n-Alkyl (68% C ₁₂ , 32% C ₁₁) dimethyl ethylbenzyl ammonium chlorides | 5% |
| Inert Ingredients: | 90% |
| | <hr/> 100% |

Ark San is a concentrate and should be diluted before using.

EPA Reg. No. 4169-2

EPA Est. No. 01674MA01

Batch Code No.

Net Contents

DANGER: KEEP OUT OF REACH OF CHILDREN. (See other precautionary and first aid statements on right side panel.)

Manufactured by

ABORN CHEMICAL INDUSTRIES, INC.

(Wholly Owned Subsidiary of M. D. Stetson Company)

168 "A" Street — South Boston, Mass. 02210

ARK SAN

Cold Disinfection

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ ounce Ark San per gallon of water for 10 minutes.

FOR ROOM FOGGING

Directions:

To minimize the danger of cross-contamination from environmental surfaces apply solution of Ark San as a fog as described below, before applying standard routine terminal cleaning and disinfecting practices. 1. Remove all human, animal and plant life from room. 2. Open closet doors and drawers. 3. Set up 34" revolving platform in center of room. 4. Mount Jet Fog-Master Tri-Jet Model 6208 or other fogging device delivering equivalent spray at 3 RPM. 5. Fill sprayer reservoir with 1.5 oz. of Ark San to 1 gallon of water to produce 1150 ppm active ingredients. 6. Set sprayer mechanism to deliver 1 gallon of solution. 7. Fog for 15 minutes for an average 2100 cu. ft. room. For different room sizes, vary spray time proportionately to ensure complete wetting of exposed surfaces. 8. Wait 2 hours before entering room after treatment. 9. Rinse fogging equipment thoroughly with clear water following use.

DANGER: Keep Out of Reach of Children. Corrosive Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before re-use. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed. Rinse empty container thoroughly with water and discard.

