


Betco 256

For Hospital & Institutional Use
Highly Concentrated Germicide Based on a Unique Blend of Quaternaries with Organic Soil Load Tolerance.
Kills HIV On Pre-Cleaned Environmental Surfaces/Objects Previously Soiled With Blood/Body Fluids.*

Cleaner/Disinfectant Deodorizer/Fungicide Virucide*

306



| | |
|---|--------|
| Copy Deryl Quaternary Ammonium Chloride | 3.75% |
| Detyl Quaternary Ammonium Chloride | 2.5% |
| Detyl Quaternary Ammonium Chloride | 2.5% |
| Allyl C- 10%, C- 40%, C- 10% | |
| Quaternary Ammonium Chloride | 1.50% |
| BEST INGREDIENTS | 27.25% |
| TOTAL | 38.25% |

KEEP OUT OF REACH OF CHILDREN
DANGER
See label panel for additional precautionary statements.
FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY.

EPA EST. NO. 4170-CH-1
EPA REG. NO. 4170-14

MAR 22 1991
4170-14

BEST AVAILABLE COPY

DESCRIPTION

Cross-contamination is of major housekeeping concern not only in hospitals, but in schools, institutions, and industry. This product is formulated for this problem area. It both cleans and disinfects effectively and is virucidal when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-contamination from environmental surfaces.

When used as directed, this product will deodorize surfaces in toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and will inhibit the growth of mold and mildew and their odors when used as directed.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Keep out of reach of children. Corrosive. Causes eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

Authorized by the USDA for use in federally inspected meat and poultry plants.

5 GALLONS

Manufactured By:
Betco Corp., Toledo, Ohio 43607

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a proven "one-step" cleaner-disinfectant or fungicide-virucide in the presence of moderate amounts of organic soil. Apply this product to walls, floors and other hard (inanimate) non-porous surfaces such as tables, chairs, counter tops, sinks and bed frames with a cloth, mop or mechanical spray device so as to thoroughly wet surfaces. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or when use solution becomes visibly dirty.

Disinfection-To disinfect hard, non-porous surfaces, add 1/2 oz. per gallon of water. Treated surfaces must remain wet for 10 minutes. At this use-level, this product is also virucidal and fungicidal.

1/2 oz. per gallon use level-The broad spectrum activity of this product has been evaluated in the presence of 5% serum and found to be effective against the following by the AOAC Use-Dilution Test.

- Pseudomonas aeruginosa
- Staphylococcus aureus
- Salmonella choleraesuis
- Brevibacterium ammoniagenes
- Klebsiella pneumoniae
- Enterobacter aerogenes
- Salmonella schottmuelleri
- Pseudomonas cepacia
- Escherichia coli
- Streptococcus faecalis
- Shigella dysenteriae

AOAC Fungicidal Test-This product is effective against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, shower and bath areas, exercise areas, dressing rooms.

Virucidal Performance-At 1/2 oz. per gallon use-level, this product was evaluated in the presence of 5% serum and found to be effective against the following viruses: Herpes Simplex Type 1 (a member of the virus family that causes infectious mononucleosis), Herpes Simplex Type 2 (the venereal disease type of the Herpes virus), Vaccinia (representative of the pox viruses), HIV-1 (AIDS virus) and Influenza A2 (the Hong Kong flu virus) on hard, non-porous inanimate environmental surfaces.

** Kills HIV on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.


CONTACT TIME: Leave surface wet for 10 minutes with 1 oz. per gallon use-solution.

CLEANING PROCEDURE: Blood and other body fluids containing HIV must be thoroughly cleaned from surfaces and objects before application of this product.

PERSONAL PROTECTION: Clean-up should always be done wearing protective latex gloves, gowns, masks and eye protection.

DISPOSAL OF INFECTIOUS MATERIAL: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

123060



BEST AVAILABLE COPY