

Sanibet™



**Disinfectant, Sanitizer, Fungicide, Deodorizer
For Hospital, Institutional, Industrial, School,
Dairy and Other Farm Use**

ACTIVE INGREDIENTS:
Didecyl dimethyl ammonium chloride 1.5%
INERT INGREDIENTS 98.5%
TOTAL 100%

**KEEP OUT OF REACH OF CHILDREN
DANGER**
See label panel for additional safety information.
FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY.



EPA REG. NO. 4170-15
EPA EST. NO. 4170-01-1

305

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with the labeling.

Apply disinfectant with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 15 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

SANITIZING OF FOOD PROCESSING EQUIPMENT AND OTHER HARD SURFACES IN FOOD CONTACT LOCATIONS

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, cutlery, glasses, and tops, counter tops, refrigerated storage and display equipment and other hard, non-porous surfaces, no potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. disinfectant in 4 gallons of water (150 ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, counters, refrigerated storage and display equipment and other stationary hard, non-porous surfaces by mop or brush. No potable water rinse is required.

Disinfectant, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz. 4 gallon solution of disinfectant. No potable water rinse is required.

At 1 oz./4 gallons, disinfectant meets the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance 1975 Recommendations of the U.S. Public Health Service in and/or up to 1.00 ppm of hardness calculated as CaCO₃ which is tested by the AOAC Germicidal and Disinfectant Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, teats, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. disinfectant in 4 gallons of warm water. No potable water rinse is required.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back in sanitizing solution, when solution becomes visibly dirty, discard and prepare fresh solution.

DISINFECTION IN HOSPITALS, NURSING HOMES AND OTHER HEALTH CARE INSTITUTIONS. For disinfecting floors, walls, counter tops, bathing areas, corridors, bed frames, tables, chairs, garbage pails and other hard non-porous surfaces.

Add 1 1/2 oz. disinfectant to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use-level, disinfectant is effective against *Pseudomonas aeruginosa*. **DISINFECTANT IN INSTITUTIONS, INDUSTRY, AND SCHOOLS.** For disinfecting floors, walls, bed frames, counter tops, tables, chairs, garbage pails, bathroom fixtures, and other hard, non-porous surfaces.

Add 2 oz. of disinfectant to 4 gallons of water. Apply to previously cleaned hard surface with mop cloth.

At 2 oz./4 gallon use-level disinfectant is effective against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Typhlophlyctis muris* (the above is foot fungus).

DISINFECTION OF PRECLEANED BARBER TOOLS. Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2-oz. gallon solution of disinfectant.

DISINFECTION OF POULTRY EQUIPMENT, ANIMAL QUARTERS AND KENNELS. Poultry brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. disinfectant in 4 gallons of water with a mop, cloth or brush. Small drains should be immersed in this solution.

Prior to disinfection, all poultry, other animals, and feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces contacted or traversed by poultry and other animals.

After disinfection, portable buildings, ramps and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Authorized by the USDA for use in federally inspected meat and poultry plants.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **DANGER**

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Possible mucous membrane damage may contraindicate the use of gastric lavage. Measure against circulatory shock, respiratory depression, and convulsion may be needed.

STORAGE AND DISPOSAL

STORE IN ORIGINAL CONTAINER ONLY

PESTICIDE DISPOSAL. Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dump of in a sanitary landfill, or by other procedures approved by the state or local authorities.

Manufactured By
Betco Corp., Toledo, Ohio

ONE GALLON

