

PM31

4170-41

ACCEPTED
AUG 19 1993
 Under Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4170-41



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. When used as directed, this germicidal cleaner demonstrates effective disinfection against *Staphylococcus aureus*, *Salmonella choleraesuis*, and *Pseudomonas aeruginosa*, thus meeting requirements for hospital use. Efficacy tests have demonstrated that this germicidal cleaner is an effective one-step cleaner/disinfectant in the presence of organic soil (5% blood serum). The product is fungicidal against Pathogenic fungi (*T. mentagrophytes*) when used as directed on hard surfaces found in the bathroom, shower stall, and locker room, or other clean, hard, non-porous surfaces commonly contacted by bare feet. **MOLD AND MILDEWSTAT**—This product prevents and controls mold and mildew and the odors they cause on hard, non-porous, inanimate surfaces. This germicidal cleaner has been designed specifically for institutional cleaning and disinfecting in nursing homes, hospitals, and institutions. **DISINFECTION, DEODORIZATION, AND CLEANING:** To disinfect hard, non-porous surfaces (such as bathroom tiles, walls, shower stalls, sinks, and bathtubs), proceed as indicated below: (1) Shake can before using. (2) Hold six to eight inches from surface to be treated. (3) Spray area until it is covered with a white foam. Allow foam to penetrate and remain wet for 10 minutes. No scrubbing necessary. (4) Wipe off with a clean cloth, mop or sponge. 360° valve allows spray in any direction. Pre-clean all surfaces to be disinfected. **KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) associated with AIDS. **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS virus) OF SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS. PERSONAL PROTECTION:** When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of disinfectant. **CONTACT TIME:** Allow surface to remain wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal. **MILDEWSTAT**—To control mold and mildew on pre-cleaned, hard, non-porous surfaces (such as floors, walls, table tops), spray the surfaces to be treated, making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth reappears. **ANTIVIRAL PERFORMANCE:** This product was evaluated and found to be effective against Herpes Simplex Type 1 (virus that causes cold sores) and Herpes Simplex Type 2 (virus that causes genital herpes) on pre-cleaned, hard, non-porous, inanimate environmental surfaces for a contact time of 10 minutes. **RESIDUAL SELF-SANITIZING ACTIVITY:** This product has been evaluated and found to have residual self-sanitizing activity in 5 minutes on pre-cleaned, hard, non-porous, inanimate environmental surfaces. This residual sanitizing activity is applicable to surfaces that are reasonably expected to become wet under normal conditions such as shower and restroom fixtures. This residual self-sanitizing activity will continue for 5 days or until the surface is rinsed or cleaned.

PRECAUTIONARY STATEMENTS
 Hazards to Humans and Domestic Animals

WARNING

Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

STORAGE AND DISPOSAL

Do not reuse empty container. Replace cap and discard container in trash. Do not incinerate or puncture.

Manufactured By:



Beteo Corp., Toledo, Ohio 43607



Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols

EPA EST. NO. 4170-OH-1
 EPA REG. NO. 4170-41

6883

GC/88

Cide-Bet
 Hospital Disinfectant
 Foaming Germicidal
 Cleaner
 Limpiador germicida espumoso
 Fresh & Clean Scent

Kills HIV on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids

* Effective against Herpes Simplex Type 1 and Type 2 (See side panel)

** Five day residual self-sanitizing activity

Authorized by the USDA for use in federally inspected meat and poultry plants

ACTIVE INGREDIENTS

n-Alkyl (C12-C15) 3-(1,3,5-triazol-2-yl) ammonium chlorides	0.1%
n-Alkyl (C12-C15) 3-(2,4,6-triazin-5-yl) dimethyl ethylbenzyl ammonium chlorides	0.1%
INERT INGREDIENTS	99.8%
TOTAL	100%

Contains Sodium Chloride

KEEP OUT OF REACH OF CHILDREN

WARNING:

See side panel for additional precautionary statements for use in meat and poultry plants only

Net Wt. 18.0 oz (510 g)

1891