



DESCRIPTION

This disinfectant toilet bowl and urinal cleaner is designed for Hospital, General Industrial and Institutional use (schools, factories, nursing homes). Its broad spectrum activity cleans, descales, deodorizes and disinfects in one easy step.

PRECAUTIONARY STATEMENTS

HAZARDOUS TO HUMANS
AND DOMESTIC ANIMALS

DANGER POISON

CORROSIVE: Keep Out of Reach of Children. May be fatal or harmful if swallowed. Do not breathe vapors or fumes. Produces chemical burns. Do not get in eyes, on skin, or on clothing. Do not use on sinks, bathtubs, porcelain or any surface other than toilet bowls & urinals.

STORAGE AND DISPOSAL

Store in original container only. Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. Drink a teaspoonful or more of milk of magnesia or cup of soapy water followed with milk or raw egg whites. **EYE CONTACT:** Wash with water for 15 minutes. Get prompt medical attention. **SKIN CONTACT:** Wipe off acid gently, flood surface with water using soap freely, then cover with moist magnesia or baking soda.

EPA EST. NO. 4170-OH-1
EPA REG. NO. 4170-52

ONE QUART



Manufactured By:
Betco Corp., Toledo, Ohio

Pull®
Hospital Type

Kills human immunodeficiency virus Type 1 (HIV-1)
(AIDS virus) on pre-cleaned environmental surfaces/
objects previously soiled with blood/body fluids.*

Disinfectant Toilet Bowl & Urinal Cleaner
Cleans, Disinfects, Deodorizes, Descales
-With Organic Soil Tolerance

ACTIVE INGREDIENTS:

Octyl decyl dimethyl
ammonium chloride0.0150%
Dioctyl dimethyl
ammonium chloride0.0075%
Didcyl dimethyl
ammonium chloride0.0075%
Alkyl (C14, 50%; C12 40%;
C10 10%) dimethyl benzyl
ammonium chloride0.0200%
Hydrogen chloride23.0000%
INERT INGREDIENTS: 76.9500%
TOTAL 100.000%

KEEP OUT OF REACH
OF CHILDREN
DANGER POISON
FOR INDUSTRIAL AND
INSTITUTIONAL USE ONLY

See side panel for additional
precautionary and practical
treatment statements.



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

TO CLEAN AND DISINFECT TOILET BOWLS AND URINALS: Remove water from bowl by forcing it over the trap with a swab mop. Saturate swab mop with cleaner/disinfectant and wet all surfaces thoroughly. It is most important to swab under rim of bowl or urinal to clean and disinfect water outlets. Allow cleaner/disinfectant to remain for at least 10 minutes until all rust, scale and stains are loosened. Rinse by flushing.

For heavily soiled areas a preliminary cleaning is required. At a dilution of one part cleaner/disinfectant to 3 parts water, this product has shown broad spectrum activity against microorganisms commonly found in toilet bowls and urinals, and has been evaluated in the presence of 5% serum and found to be effective against the following organisms by the AOAC Use-Dilution test:

PSEUDOMONAS AERUGINOSA
STAPHYLOCOCCUS AUREUS
SALMONELLA CHOLERAESUIS

*In health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS). **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Allow surface to remain wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

BEST AVAILABLE COPY