



ACTIVE INGREDIENTS	
Copper, Expressed as Metal	4.00
Derived from Cuprous and Cupric Oxide	
Sulfur	84.0
	110
INERT INGREDIENTS	
TOTAL	100

### DIRECTIONS

FOR THE CONTROL OF LEAF SPOT and LEAF HOPPER  
For the Bunch or Runner types, make first application during the first ten days of July. On Spanish, dusting should be started about ten days earlier, following with two or three applications at as nearly fourteen day intervals as weather conditions will permit. The rate of application per acre should range from 15 to 20 lbs. per acre totaling approximately 70 lbs. per acre in four applications or 50 lbs. in three. Should an application be washed off within a few hours, the treatment should be repeated as soon as weather conditions will permit.

### CAUTION

Do not breathe dust. Do not get in eyes, in skin or on clothing. Dispose of wastes by burying in open pits away from water supplies. Containers should be disposed of by burying with wastes or by burning. (Keep out of smoke)

**NOTICE OF WARRANTY.** The Borden Chemical Company, Safety Division, warrants only that the contents of the container correspond to the analysis printed on the container label. THERE IS NO WARRANTY OF MERCHANTABILITY AND THERE ARE NO WARRANTIES THAT EXTEND BEYOND SUCH ANALYSIS. Buyer assumes all risks of use of such material in accordance with directions on the label.

**CAUTION:** KEEP OUT OF THE REACH OF CHILDREN

See "Appendix" p. 10.

Distributed by THE BORDEN CHEMICAL COMPANY

\* \* \* \* \*

USDA-ARS • 2016 • 9781614762414