



**1-67**

4-26-67  
4185-94

ACTIVE INGREDIENTS:

Malathion\*

4.0%

TOTAL 100.0%

INERT INGREDIENTS

\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

96.0%

**DIRECTIONS**

**CABBAGE, KALE, COLLARDS, MUSTARD, TURNIP and BROCCOLI:** To control Aphids, Imported Cabbage Worm and Cabbage Looper, dust at the rate of 30 to 40 lbs. per acre, repeating when necessary. Make no application within 7 days of harvest to Cabbage, Kale Collards. Mustard and Turnip; within 3 days of harvest of Broccoli.

**BEANS:** To control Mexican Bean Beetles, Leafhopper and Spider Mites, dust at the rate of 30 to 35 lbs. per acre, repeating when necessary. Make no application within 24 hours of harvest.

**PEAS:** To control Pea Aphid, dust at the rate of 25 to 35 lbs. per acre, repeating when necessary. Make no application within 3 days of harvest.

**TOMATO:** To control Spider Mites and Aphids, dust at the rate of 35 to 45 lbs. per acre, repeating when necessary. Make no application later than 3 days before harvest.

**POTATO:** To control Aphids, dust at the rate of 30 to 35 lbs. per acre, repeating as necessary. Make no application later than 3 days before harvest.

**CUCUMBER, SQUASH and MELON:** To control Aphids and Spider Mites, dust at the rate of 30 to 35 lbs. per acre, repeating when necessary. Do not dust unless plants are dry and make no application within 3 days of picking.

**PEPPERS: AND EGGPLANTS:** To control Aphids and Spider Mites, dust at the rate of 20 to 25 lbs. per acre, repeating when necessary. Make no application within 3 days of harvest.

**ORNAMENTALS:** For the control of the following: Scuffy Scale and Magnolia Scale, Pine Leaf Scale, Eucalyptus Scale, Scale, European Pine Shoot Moth, Leaf Caterpillar, Minor, Lace Bugs, Grasshoppers, White Fly, Scale Crawlers, Aphids, and Red Spider. Make thorough for when necessary. Applications for scale control should be at crawler stage. Do not use on certain ferns including Boston, as on some species of Crassula.

**CAUTION**

Harmful if swallowed. Avoid prolonged breathing of dust. Contact with skin; wash thoroughly after using. Avoid contact with eyes. Do not breathe dust. Do not get in eyes, on skin, is poisonous to fish. Keep out of lakes, streams, and ponds. Conditions favor drift of dust from areas treated. Apply this label.

Dispose of wastes by burying in non-crop lands away from crops. Should be disposed of by burying with wastes, or by burning.

**NOTICE OF WARRANTY:** The Borden Chemical Company, Smith-Douglass Division, only that the contents of the container correspond to the label, label, or tag. THERE IS NO WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE. NO WARRANTIES THAT EXTEND BEYOND SUCH ANALYSIS. Use, storage or handling, whether in accordance with directions on this label.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

See other cautions on this label

Distributed by THE BORDEN CHEMICAL CO.

SMITH-DOUGLASS DIVISION • Norfolk, Va.

ACTIVE INGREDIENTS:	
Malathion*	4.0%
	TOTAL 100.0%
INERT INGREDIENTS 96.0%	
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate	

CTIONS

RNIP and BROCCOLI: To control Aphids, Im-  
dust at the rate of 30 to 40 lbs. per acre, re-  
n within 7 days of harvest to Cabbage, Kale  
s of harvest of Broccoli.

eanhopper and Spider Mites, dust at the rate  
ecessary. Make no application within 24 hours

te of 25 to 35 lbs. per acre, repeating when  
of harvest.

ids, dust at the rate of 35 to 45 lbs. per  
plication later than 3 days before harvest.

ate of 30 to 35 lbs. per acre, repeating as  
ays before harvest.

ol Aphids and Spider Mites, dust at the rate  
necessary. Do not dust unless plants are dry  
king.

ids and Spider Mites, dust at the rate of 20  
ssary. Make no application within 3 days of

ORNAMENTALS: For the control of the following: Scurfy Scale, Florida Red Scale, Azalea  
and Magnolia Scale, Pine Leaf Scale, Eucalyptus Scale, Soft Brown Scale, Monterey  
Scale, European Pine Shoot Moth, Tent Caterpillar, Black Leaf Miner and Boxwood Leaf  
Miner, Lace Bugs, Barkweevil, White Fly, Japanese Beetle, Black  
Scale Crawlers, Aphids, and Red Spider. Make thorough full cover applications, repeating  
when necessary. Applications for scale control should be made when scales are in the  
crawler stage. Do not use on certain ferns including Boston, Maidenhair and Pteris, as well  
as on some species of Crassula.

CAUTION

Harmful if swallowed. Avoid prolonged breathing of dust. Avoid prolonged or repeated  
contact with skin; wash thoroughly after using. Avoid contamination of feed and food-  
stuffs. Do not breathe dust. Do not get in eyes, on skin, or on clothing. This product  
is poisonous to fish. Keep out of lakes, streams, and ponds. Do not apply when weather  
conditions favor drift of dust from areas treated. Apply this product only as specified on  
this label.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers  
should be disposed of by burying with wastes, or by burning (Keep out of smoke).

NOTICE OF WARRANTY: The Borden Chemical Company, Smith-Douglass Division, warrants  
only that the contents of the container correspond to the analysis printed on the con-  
tainer, label, or tag. THERE IS NO WARRANTY OF MERCHANTABILITY AND THERE ARE  
NO WARRANTIES THAT EXTEND BEYOND SUCH ANALYSIS. Buyer assumes all risks of  
use, storage or handling, whether in accordance with directions or not.

AUTION: KEEP OUT OF REACH OF CHILDREN

See other cautions on this label