



ACTIVE INGREDIENTS:	
Toxaphene-Technical Chlorinated Camphene	
(Chlorine Content 67-69%)	20.0%
INERT INGREDIENTS	80.0%
TOTAL	100.0%

DIRECTIONS

CABBAGE, BROCCOLI, LETTUCE AND BRUSSELS SPROUTS: To control Cabbage Loopers, apply at the rate of 10 lbs. per acre. For Cutworms (soil), apply 20 lbs. per acre to soil surface. For Cabbage and Head Lettuce, apply up to 7 days before harvest, if the outer leaves are stripped at harvest. If the outer leaves are to be left on at harvest, do not apply after the heads begin to form. For Leaf Lettuce, do not apply after seedling stage. For Broccoli and Brussels Sprouts, do not apply after the edible parts begin to form. On all the above, do not apply this product if leaves, stems or heads are to be fed to dairy animals or meat animals.

BOLLARDS AND KALE: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface Feeding Cutworms, apply at the rate of 10 to 20 lbs. per acre. Do not apply more than 4 times nor more than 3 times after thinning. Do not apply within 14 days of harvest when 10 lbs. per acre is used, do not apply within 21 days of harvest when 20 lbs. per acre is used.

SPINACH: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface Feeding Cutworms, apply only once at the rate of 10 lbs. per acre. Do not apply within 21 days of harvest.

BEANS: To control Cutworms (soil), apply 10 lbs. per acre on soil in late afternoon. Do not apply after pods begin to form.

TOMATOES: To control Tomato Hornworm, Tomato Fruit Worm, Serpentine Leaf Miner, Leaf Beetles, Cutworms, Cabbage Looper, Russet Mites and Colorado Potato Beetles, apply 10 to 20 lbs. per acre as required. Do not apply within 5 days of harvest.

SOYBEANS: To control Velvet Bean Caterpillars, Corn Earworms, Bean Leaf Beetles, Army Worms and related insects, apply at the rate of from 20 to 30 lbs. to the acre. Do not feed treated forage to dairy animals or animals being finished for slaughter.

PEANUTS: To control Fall Army Worm and Velvet Bean Caterpillar, apply at the rate of 7½ to 10 lbs. per acre. Do not feed foliage to dairy animals or animals being finished for slaughter.

CORN: To control Army Worm and Fall Army Worm, apply at the rate of 20 lbs. per acre. Do not feed treated plants to dairy or meat animals.

SORGHUM: To control Army Worm and Fall Army Worm, apply at the rate of 10 lbs. per acre. Do not feed treated plants or ensilage from treated plants to dairy or meat animals. Do not use on Sorghum to be sold commercially or to be shipped interstate. Do not apply within 40 days of harvest or apply after heads have begun to form.

WHEAT, BARLEY, OATS AND RYE: To control Climbing and Surface Feeding Cutworms, Army Worms, Grasshoppers and Chinch Bugs, apply at the rate of 10 lbs. per acre. Do not apply more than once after heads have begun to form. Do not apply within one week of harvest of Wheat, Rye and Oats within 2 weeks of harvest of barley. Do not pasture dairy animals or meat animals in treated fields.

Destroy bags by punching holes in them and bury or burn. Keep out of smoke.

CAUTION

Harmful if swallowed. Do not breathe dust or spray. Do not get on skin, hair or clothing. Wash thoroughly after using. Avoid contact with eyes and mucous membranes. Do not use on cucurbits. Do not use on feed or forage.

This product is poisonous to fish and wildlife. Keep out of streams and ponds. Do not apply when weather conditions favor drift. Do not use on treated crops. Apply product only as specified on this label.

NOTICE OF WARRANTY: The Borden Chemical Company, Inc. warrants only that the contents of the container conform to the label. There is no warranty of merchantability and there are no warranties that extend beyond such analysis. Buyer assumes all risk of use, storage or handling, whether in accordance with directions or not.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

See other cautions on this label

Distributed by THE BORDEN CHEMICAL CO.