



ACTIVE INGREDIENT: Captan* 7.5%
INERT INGREDIENTS 92.5%
TOTAL 100.0%

*N-Trichloromethylthio-4-cyclohexene-1, 2-dicarboximide

DIRECTIONS

STRAWBERRIES: To control Gray Mold or Botrytis Blight — Apply at the rate of 35 to 40 lbs. per acre at one week intervals from the time plants start to grow actively in the spring.

POTATOES (SEED PIECES): To control Seed Rot, Damping off, Verticillium Wilt and Brown Eye Disease — Use at the rate of 10 ozs. per bushel cut seed pieces; (1 lb. per 100 lbs. cut seed pieces). Apply in convenient container or by dust attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days, the treated pieces should be stored for 2 to 3 days in open crates before bagging.

CAUTION

Avoid inhaling dust. Wear a respirator approved by the U. S. Department of Agriculture for Captan while handling. Avoid prolonged or repeated contact with skin. Avoid contact with eyes skin or clothing. Wash thoroughly after using. Avoid contamination of feed and foodstuffs.

This product is poisonous to fish. Keep out of lakes, streams, and ponds. Do not apply when weather conditions favor drift of dust from areas treated. Apply this product only as specified on this label.

Store in cool, dry place. Protect from excessive heat. Burn bag immediately when empty. Stay out of smoke.

NOTICE OF WARRANTY: The Borden Chemical Company, Smith Douglass Division, warrants only that the contents of the container correspond to the analysis printed on the container, label, or tag: THERE IS NO WARRANTY OF MERCHANTABILITY AND THERE ARE NO WARRANTIES THAT EXTEND BEYOND SUCH ANALYSIS. Buyer assumes all risks of use, storage or handling, whether in accordance with directions or not.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See other cautions on this label

Distributed by THE BORDEN CHEMICAL CO.

SMITH-DOUGLASS DIVISION • Norfolk Va