




44-299
05-411

ACTIVE INGREDIENTS	
Carbaryl (1 naphthyl N methylcarbamate)	5.00%
Parathion (O O diethyl O p nitrophenyl thiophosphate)	1.00%
INERT INGREDIENTS	
TOTAL	94.00%
	100.00%

Trade Mark of Union Carbide Corporation for 1 naphthyl N methylcarbamate.

DIRECTIONS FOR USE

HEAD ENTIRE APL USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND WARNINGS

TOBACCO

Hand Dusting Application: For budworms and tobacco flea beetles on individual plants, apply 5 to 8 pounds per acre. Heavier applications may cause plant injury. Rotary dusting is recommended.

Broadcast Application: For hornworms, budworms, aphids, flea beetles, suckfly, grasshoppers and Japanese beetle, apply 10 to 40 pounds per acre depending on size of the plants. Use the highest rate only on large plants. Full coverage is essential. Best treatment when worms are small and repeat as needed. Heavier applications may cause plant injury.

For late applications are necessary, treat immediately after pruning. Do not apply within 5 days of pruning or 15 days of cutting. Avoid getting wet with plant juices when cutting. Do not use phytotoxicity may occur on tender foliage in the presence of rain or high humidity at several days after application.

VEGETABLE CROPS

IN ALL VEGETABLE OR FRUIT CROPS LISTED BELOW USE AT THE RATE OF 5 TO 10 POUNDS PER ACRE. OBSERVE ALL SPECIAL USE PRECAUTIONS PERTAINING TO EACH CROP.

CORN POTATOES: To control Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. Do not apply within 5 days of harvest.

TOMATOES: To control Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bug, and stink bugs. Do not apply to tomatoes within 15 days of harvest.

CABBAGE: To control flea beetle, harlequin bug and leafhoppers, armyworms, cabbage rootworms, cutworms. For small cabbage weevils apply 5 to 10 pounds per acre. Make no more than 2 applications per season. Do not apply within 15 days of harvest.

Apply to control flea beetle, harlequin bug, leafhoppers, armyworms, cabbage rootworms, cutworms. Do not apply within 15 days of harvest.

PEPPERS: To control Mexican bean beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bug. Do not apply within 15 days of harvest.

FIELD & SWEET CORN: To control European corn borer, fall armyworm, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bug, and stink bugs. Do not apply within 15 days of harvest.

Sweet corn application to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Repeat as necessary for insects attacking silks and ears, apply at 2 to 3 day intervals when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation. Do not apply within 12 days of harvest. Timing and good coverage are absolutely essential for effective control.

PEPPERS: To control Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. Do not apply within 15 days of harvest.

SOYBEANS: To control blister beetles and Mexican bean beetle, alfalfa caterpillar, bean leaf beetle, cucumber beetles, green cloverworm, Japanese beetle, leafhoppers, three cornered alfalfa hopper, thrips and velvet bean caterpillar. Do not apply within 15 days of harvest. Some leaf injury may occur if applications are made when foliage is wet.

WARNING

MAKING: PARSLEY, WALLOWED, impaled, or absorbed through skin. Rapidly absorbed through skin. Do not get on eyes. If on face, wash immediately with soap and water. Wear a mask or respirator if a face mask is used by the U.S. Department of Agriculture for parathion protection. Do not apply when weather conditions favor drift of dust from areas treated. Keep a complete record of use, date, and area, and only where there may be a danger of drift, avoid areas that may be hazardous to other crops. Do not use in areas where there is a danger of drift. Wash hands and face thoroughly with soap and water before eating, drinking, or smoking. Wash a contaminated cloth with soap and hot water before reuse.

This product is poisonous to fish and wildlife. Keep out of water, streams, and ponds. Do not use in areas where there is a danger of drift.

Do not use in areas where there is a danger of drift. Do not use in areas where there is a danger of drift.

Do not use in areas where there is a danger of drift. Do not use in areas where there is a danger of drift.

Do not use in areas where there is a danger of drift. Do not use in areas where there is a danger of drift.

Do not use in areas where there is a danger of drift. Do not use in areas where there is a danger of drift.

Do not use in areas where there is a danger of drift. Do not use in areas where there is a danger of drift.

Do not use in areas where there is a danger of drift. Do not use in areas where there is a danger of drift.

Do not use in areas where there is a danger of drift. Do not use in areas where there is a danger of drift.

WARNING: KEEP OUT OF THE REACH OF CHILDREN

See Other Warnings On This Label

Distributed by SMITH-DOUGLASS
DIVISION OF BORDEN CHEMICAL, BORDEN INC.
Norfolk, Va. 23501