



RESTRICTED USE PESTICIDES
For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's Certification

121
P. 12
415-112

SEVIN* - PARATHION 5-1 DUST

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methylcarbamate)	5.00%
Parathion (O, O diethyl O p nitrophenyl thiophosphate)	1.00%
INERT INGREDIENTS	94.00%
TOTAL	100.00%

*Trade Mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

READ ENTIRE LABEL
CAUTIONS

DIRECTIONS FOR USE
ACCEPTED
JUL 2 1979
Under the Federal Insecticide,
Fungicide and Rodenticide Act
as amended by the pesticide
control act of 1972
4185-132

VEGETABLE CROPS

IN ALL VEGETABLE OR FRUIT CROPS LISTED BELOW USE AT THE RATE OF 25 TO 30 LBS. PER ACRE. OBSERVE ALL SPECIAL USE PRECAUTIONS PERTAINING TO SPECIFIC CROPS.
POTATOES: To control Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. Do not apply within 5 days of harvest.
TOMATOES: To control Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. Do not apply to tomatoes within 10 days of harvest.
CABBAGE: To control flea beetles, harlequin bug and leafhoppers, armyworms, cabbage caterpillars, corn earworm. For small cabbage, apply on a 5-7 day schedule. Make no more than 2 applications per season. Do not apply within 10 days of harvest.
BROCCOLI: To control flea beetles, harlequin bug, leafhoppers, armyworms, cabbage caterpillars and corn earworm. Do not apply within 7 days of harvest.
BEAN: (Green beans, snap beans) To control Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, bean cutworm, armyworm, corn earworm, stink bugs and tarnished plant bug. Do not apply within 15 days of harvest.
PEAS & SWEET CORN: To control corn rootworm adults, European corn borer, fall armyworm, flea beetles, leafhoppers, asp beetles and leafhoppers. For areas in which rootworm damage is a life long problem, apply to entire plant. To control corn earworm on

sweet corn, application to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation. Do not apply within 12 days of harvest. Timing and good coverage are absolutely essential for effective control.
PEPPERS: To control Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. Do not apply within 15 days of harvest.
SUBBEANS: To control blister beetles and Mexican bean beetle, alfalfa caterpillar, bean leaf beetle, cucumber beetles, green cloverworm, Japanese beetle, leafhoppers, three cornered alfalfa hopper, thrips and velvet bean caterpillar. Do not apply within 20 days of harvest. Some leaf injury may occur if applications are made when tender foliage is wet.

WARNING

MAY BE FATAL IF SWALLOWED, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for parathion protection. Do not apply when weather conditions favor drift of dust from areas treated. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Varied areas should not be reentered until drifting insecticides and volatile residues have dissipated. Workers entering a treated field within 5 days of an application should wear protective clothing. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and hot water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. This product is poisonous to fish and wildlife. Keep out of lakes, streams and ponds. Apply this product only as specified on this label. Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by burying with wastes. For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.
NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but seller makes NO WARRANTY OF MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or any other warranty express or implied except as herein stated. Sellers liability is limited to the purchase price of the product and this shall be the buyers sole remedy. Buyer assumes the risk of use of this product contrary to label instructions, carelessly or under conditions not intended by or not reasonably foreseeable to seller.

WARNING: KEEP OUT OF THE REACH OF CHILDREN

See Other Warnings On This Label
Distributed by SMITH DOUGLASS
DIVISION OF BORDEN CHEMICAL CORPORATION
Norfolk, Va. 23511
BEST AVAILABLE COPY

EPH 22-481-72
EPH 22-481-72