

1 (of 2)

SMITH-DOUGLASS

PHOSDRIN 2% DUST

ACTIVE INGREDIENTS:

Alpha Isomer of 2-Carboethoxy-1-methylvinyl Dimethyl Phosphate.....	1.2%**
Related Compounds.....	0.8%**

INERT INGREDIENTS:	98.0%
Total	100.0%

*Trademark

**Equivalent to 2.0% w Phosdrin Insecticide



DANGER

POISON

18 point type

DANGER : KEEP OUT OF REACH OF CHILDREN

12 point type

See antidote and other warnings on this label



Distributed by

The Borden Chemical Company
Smith-Douglass Division

Norfolk, Virginia

USDA Reg. No. 4185-141

DIRECTIONS FOR USE



Use this product only for recommended purposes and at recommended dosages.

CABBAGE AND BROCCOLI: For Aphids, apply at the rate of 8 to 12 lbs. per acre. For Cabbage Looper, Imported Cabbage Worm, Leafhoppers, Salt-Marsh Caterpillar, Mites, Cutworm (climbing), Dipterous Leafminer (adult), and Lygus Bugs, apply at the rate of 15 to 25 lbs. per acre. Repeat applications when necessary. Do not apply within 1 day of harvest.

KALE, COLLARDS, TURNIP TOPS, MUSTARD GREENS AND BRUSSELS SPROUTS: For Aphids, apply at the rate of 8 to 12 lbs. per acre. For Cabbage Looper, Imported Cabbage Worm, Leafhoppers, Salt-Marsh Caterpillar, Mites, Cutworm (climbing), Dipterous Leafminer (adult), False Chinch Bugs, and Lygus Bugs, apply at the rate of 15 to 25 lbs. per acre. Repeat applications when necessary. Do not apply within 3 days of harvest.

SPINACH: For Aphids, apply at the rate of 8 to 12 lbs. per acre. For Cabbage Looper, Imported Cabbage Worm, Leafhoppers, Mites, Dipterous Leafminer (adult), Cutworm (climbing), Salt-Marsh Caterpillar and False Chinch Bug, apply at the rate of 15 to 25 lbs. per acre. Repeat applications when necessary. Do not apply within 4 days of harvest.

LETTUCE: For Aphids, apply at the rate of 8 to 12 lbs. per acre. For Corn Earworm, Cutworm (climbing), Dipterous Leafminer (adult), Cabbage Looper, Imported Cabbage Worm, Leafhoppers, Mites, Lygus Bugs, Salt-Marsh Caterpillar and False Chinch Bugs, apply at the rate of 15 to 25 lbs. per acre. Repeat applications when necessary. Do not apply within 2 days of harvest.

USE PRECAUTIONS

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Unprotected humans and animals should be kept out of treated fields on the day of treatment. Keep out of the reach of children and domestic animals. Do not contaminate feed or feed products.

This product is poisonous to fish and wildlife. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label.

2(cf2)

WARNINGS

MAY BE FATAL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. Do not swallow or get in eyes, on skin, or on clothing. Wear clean natural rubber gloves and clean protective clothing and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Phosdrin insecticide protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least ten minutes and get medical attention. In case of spillage on person or clothing, immediately remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse.

NOTICE OF WARRANTY: The Borden Chemical Company, Smith-Douglass Division, warrants only that the contents of the container correspond to the analysis printed on the container, label or tag: THERE IS NO WARRANTY OF MERCHANTABILITY AND THERE ARE NO WARRANTIES THAT EXTEND BEYOND SUCH ANALYSIS. Buyer assumes all risks of use, storage or handling, whether in accordance with directions or not.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by burying with wastes or by burning (Keep out of smoke).

NOTE TO PHYSICIAN: Phosdrin is a cholinesterase inhibitor and treatment of severe poisoning should include the use of atropine.

