



For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's Certification.

ALC C. T. E. D
JUL 3 1979
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 4185-152

Feb 1979 Parathion 2% Dust

ACTIVE INGREDIENT:
Parathion (O, O-diethyl O-p-nitrophenyl thiophosphate)
INERT INGREDIENTS: - TOTAL

25
95
100

DIRECTIONS FOR USE

DIRECTIONS FOR USE
READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS: AND WITH APPLICABLE FEDERAL AND STATE REGULATIONS. USE 12 to 20 lbs. per acre unless otherwise specified. This product is suitable for application through ground dusters, airplane or hand equipment.
APHIDS — CABBAGE, CAULIFLOWER, AND BRUSSELS SPROUTS 15 to 20 lbs. to the acre. Not more than two applications and no application after the heads start to form or within 21 days of harvest, whichever is the earlier.
MEXICAN BLAY BEETLE — BEANS 20 to 25 lbs. to the acre. No application within 21 days before harvest.
APHIDS — PEAS 20 to 25 lbs. to the acre. Not more than two applications during the growing season and no application within 10 days of harvest time or after bloom, whichever is the earlier.
APHIDS — TOMATOES 20 to 25 lbs. to the acre. Not more than two applications and no application within 10 days of harvest.
PED SPIDER — STRAWBERRIES 25 to 30 lbs. to the acre. Not more than two applications during growing season and no application after fruit sets or 10 days before harvest.
FALL ARMY WORMS — YOUNG KALE, COLLARDS AND CABBAGE 15 to 20 lbs. to the acre. Do not apply within 7 days of cutting time in case of Kale or Collards. No application after heads have started to form in case of Cabbage.
APHIDS, RED SPIDER MITES, LEAFHOPPERS — APPLES 17 to 25 lbs. per acre. Do not apply within 14 days of harvest.
BLOSSOM WEEVILS — BLUEBERRIES 15 to 25 lbs. per acre when leaf buds show green and 7 to 9 days later. Apply only before fruit sets or after harvest.
APHIDS, LEAFMINER, THRIPS, SPIDER MITES — CELERY 30 lbs. to the acre. Do not apply within 30 days of harvest.
THRIPS — ONIONS 12 to 17 lbs. per acre. Not more than two applications per season. Do not apply within 15 days of harvest.
CUCUMBER BEETLE, MELON WORM, PICKLE WORM, SERPENTINE LEAF MINER, APHIDS, RED SPIDER MITES, THRIPS — CUCUMBER, SQUASH, MELONS, PEAS, BEANS, ETC. 15 to 20 lbs. to the acre. Apply to Cucumbers before vineing or within 15 days of harvest. Do not apply to Beans within 15 days of harvest. Do not apply to Melons within 7 days of harvest.
PEACH ACHID — PEACHES 30 to 35 lbs. per acre. Do not apply to peaches that are still on the tree. Apply cultural Experiment Station for proper timing of application. Do not apply to peaches that are still to open. Do not feed sprays to persons that are in contact with sprays to day or as a precautionary measure. Do not spray on children.
ACHID AND SERPENTINE LEAF MINER — PEPPER 15 lbs. to the acre. Do not apply within 15 days of harvest.
RED SPIDER MITES, ACHID, COLORADO POTATO BEETLE, BEANS, ETC. 15 to 20 lbs. to the acre. Apply to Beans before vineing or within 15 days of harvest. Do not apply to Beans within 15 days of harvest.
SPINACH — SPINACH 12 to 15 lbs. to the acre. Do not apply within 10 days of harvest.
WALKER ACHID — WALKER 15 to 20 lbs. to the acre. Do not apply within 10 days of harvest.

LOTON AFRID. PED SPIDERS. LEAFHOPPER — LOTON. Do not apply more than 1 or 2 times during the season. Do not apply within 5 days before harvest or within 15 days of export. Do not use on leafhoppers or on a mass basis. Limited for spraying.

SPIDERS (AFRID). SPIDERS. ME AND LON. HOPPER — LOTON. Do not apply more than 1 or 2 times during the season. Do not apply within 5 days before harvest or within 15 days of export. Do not use on leafhoppers or on a mass basis. Limited for spraying.

WHEAT CATS. HOPPER. 17 to 19. Do not apply more than 1 or 2 times during the season. Do not apply within 5 days before harvest or within 15 days of export. Do not use on leafhoppers or on a mass basis. Limited for spraying.

WARNING

May be fatal if swallowed. Do not swallow. Wash thoroughly through skin. Do not breathe dust. Rinse thoroughly with water. If in eyes, flush with water for 15 minutes. Do not use eye drops. If on face, wash thoroughly with water. Do not use eye drops. Wear a mask and gloves. Do not use in confined spaces. Do not use in areas of fire. Do not use in areas where there may be flammable vapors. Do not use in areas where there may be flammable liquids. Do not use in areas where there may be flammable gases. Do not use in areas where there may be flammable dusts. Do not use in areas where there may be flammable fumes. Do not use in areas where there may be flammable mists. Do not use in areas where there may be flammable aerosols. Do not use in areas where there may be flammable vapors. Do not use in areas where there may be flammable liquids. Do not use in areas where there may be flammable gases. Do not use in areas where there may be flammable dusts. Do not use in areas where there may be flammable fumes. Do not use in areas where there may be flammable mists. Do not use in areas where there may be flammable aerosols.

DANGER



POISON

[illegible]

the 1990s, the number of people in the world who are under 15 years of age is expected to increase from 1.1 billion to 1.5 billion. The number of people aged 65 and over is expected to increase from 250 million to 450 million. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion.

WA
BEST DOCUMENT AVAILABLE

WARNING: KEEP OUT OF REACH OF CHILDREN

See antidote and other warnings on this label

1. *Author*: Mary Ellen Dougherty