

DIRECTIONS FOR USE

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

ASPARAGUS: Use 20 - 40 pounds per acre for asparagus beetle on seedlings or spears. Use up to one day of harvest.

BEANS: (Green beans, lima beans, snap beans, cowpeas and blackeyed peas). Use 10 lbs. per acre for Mexican bean beetle; use 20 lbs. per acre for bean leaf beetle, cucumber beetles, flea beetles, Western bean cutworms; use 20 to 30 lbs. per acre for armyworms, corn earworm, stink bugs and tarnished plant bug. Use up to the day of harvest.

CORN: Use 20 to 40 lbs. per acre for corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles and leafhoppers. For larvae in whorl (Budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Use up to the day of harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI: Use 10 to 20 lbs. per acre for flea beetles and harlequin bug. Use 20 to 40 lbs. per acre for armyworms, cabbage caterpillars and corn earworm. Use 20 to 40 lbs. per acre for small cabbage loopers in a 5 to 7 day schedule. Use up to 3 days of harvest.

HEAD LETTUCE AND CARROTS: Use 10 to 20 lbs. per acre for flea beetles, harlequin bug and leafhoppers. To control six-spotted leafhopper use 20 to 30 lbs. per acre in a 5 to 7 day schedule. Use 20 to 40 lbs. per acre for armyworms, cabbage worms and loopers, corn earworm, spittlebugs, tarnished plant bugs and stink bugs. For optimum control of corn earworm after lettuce heads begin to form use 40 lbs. per acre in a 5 to 7 day schedule. Use 20 to 40 lbs. per acre for small cabbage looper in a 5 to 7 day schedule. Do not use within 3 days of harvest on lettuce; use up to day of harvest on carrots.

CUCUMBER, MELONS, PUMPKIN AND SQUASH: Use 10 to 20 lbs. per acre for pickleworm and melonworm; 20 lbs. per acre for cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida. Use up to the day of harvest.

POTATO, TOMATO, EGGPLANT AND PEPPER: Use 10 to 20 lbs. per acre for Colorado potato beetle, flea beetles and leafhoppers. Use 20 to 40 lbs. per acre for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. Use up to the day of harvest.



OKRA: Use 20 to 40 lbs. per acre for corn earworm and stink bugs. Repeat at 5 to 7 day intervals or as necessary. Use up to day of harvest.

BLUEBERRIES: Use 30 to 40 lbs. per acre for blueberry maggot. Apply 3 weeks before harvest and repeat 10 days later or as necessary. Use up to the day of harvest. Use 30 to 40 lbs. per acre for cherry and cranberry fruitworm, European fruit lecanium and Japanese beetle. Use up to the day of harvest.

STRAWBERRIES: Use 20 to 40 lbs. per acre for meadow spittlebug, strawberry leaf roller and strawberry weevil. Do not apply within 1 day of harvest.

ALFALFA, CLOVERS, COWPEAS, PEANUTS AND SOYBEANS: Use 10 to 20 lbs. per acre for blister beetles and Mexican bean beetle. Use 20 lbs. per acre for alfalfa caterpillar, bean leaf beetle, cucumber beetles, green cloverworm, Japanese beetle, leafhoppers, three cornered alfalfa hopper, thrips, and velvet bean caterpillar. Use 20 to 30 lbs. per acre for armyworm, corn earworm, stink bugs and webworms. Some leaf injury may occur if applications are made when tender foliage is wet. Use up to the day of harvest.

GRAPES: Grape Leaf Folder - 1st Brood: 20 to 25 lbs. per acre just before first brood larvae emerge from the first rolls. 2nd and 3rd Broods: 20 to 40 pounds per acre. May be applied up to harvest. Grape Leafhopper -- 20 pounds per acre. Thorough coverage is essential. This application may be expected to control DDT-resistant leafhoppers.

COTTON: Early-Season -- Thrips, Cotton Leafhopper -- 15 to 20 lbs. per acre as soon as insects appear. Mid and Late-Season -- Boll Weevil, Boll Worm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator, Aphids -- 40 lbs. per acre. Apply when insects appear and repeat at 5 to 7 day intervals. Aphid populations will usually be repressed when this insecticide is used on a regular schedule of treatments. Do not graze treated fields or feed trash to dairy or meat animals.

FILBERTS: Filbert Aphid, Filbert Leaf Roller -- 80 lbs. per acre. Apply when Leaf Roller eggs are hatching. Filbertworm -- 80 lbs. per acre. Apply when adult moths first appear and repeat three to four weeks later. Do not apply within 60 days of harvest.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-Facing Insects, Periodicals Cicada -- 70 to 80 lbs. per acre. Do not apply within 1 day of harvest.

FORAGE, GRASSES AND PASTURE: Use 20 to 30 lbs. per acre for armyworms, stink bugs and thrips. Use up to the day of harvest.

SORGHUMS: (including ^{Sorghum} Milo and Grain Sorghum): Use 20 to 40 lbs. per acre for armyworms, corn earworms, stink bugs and webworms. For optimum insect control on grain sorghum, direct dust into forming heads. Use 30 lbs. per acre for sorghum midge. Treatment for sorghum midge control should be made 3 to 4 days after heads have emerged from boot. Do not use closer than 21 days of harvest.



TOBACCO: In Plant Beds: For Tobacco Flea Beetle, apply 1 to 2 pounds per 100 square yards when plants appear. Repeat as needed at 10 to 14 day intervals and just before transplanting.

TOBACCO: In Fields: Foliage application for control of tobacco flea beetles and ~~hornworms~~, use 15 to 25 pounds per acre depending on the size of the plants. Good coverage is essential. Heavier application may cause plant injury.

NOTE: If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

WARNING

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid contact with skin. Skin contact may be harmful. Avoid contact. Wear regular long sleeved work clothing. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. Do not get in eyes, on skin, or on clothing.

Note to Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Keep out of any body of water. Fish may be killed. For protection of honeybees, avoid use, if possible, during periods when honeybees are visiting the crop or neighboring crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance (beyond bee flight range) until one week after application.

Do not reuse empty container. Punch holes in bags, and bury or burn (Keep out of smoke or fumes).

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but seller makes NO WARRANTY OF MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or any other warranty, express or implied, except as herein stated. Seller's liability is limited to the purchase price of the product and this shall be the buyers only remedy. Buyer assumes the risk of use of this product contrary to label instructions, carelessly, or under conditions not intended by or not reasonably foreseeable to seller.

