

SEVIN*—SULFUR 5-75 DUST

ACCEPTED

JUL 27 1967

UNION CARBIDE
FUNGICIDE ACT
FOR REGISTERED
ED CHLORIDE
TO ATTENTION
4185760 SUBJECT

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)
Sulfur

INERT INGREDIENTS:

TOTAL

5.00%
75.00%
20.00%

100.00%

*Trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate.

DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

GRAPES: GRAPE LEAF FOLDER: 1st Brood: 20 to 25 lbs. per acre just before first brood larvae emerge from the first rolls. 2nd and 3rd Broods: 20 to 40 pounds per acre. May be applied up to harvest. Grape Leafhopper — 20 pounds per acre. Thorough coverage is essential. This application may be expected to control DDT-resistant leafhoppers.

BEANS: MEXICAN BEAN BEETLE: 10 pounds per acre. Leafhoppers — 20 pounds per acre. Lygus, Stink Bugs — 40 pounds per acre. May be used up to harvest. Apply thoroughly at first signs of insect infestation. When necessary repeat every 7 to 10 days. Do not feed treated vines to meat or dairy animals.

CORN: CORN EARWORM: 30 to 40 lbs. per acre. Apply at 3-day intervals starting when silks first appear. Make 4 to 5 applications until silks dry or turn brown. May be applied up to harvest on sweet corn to be used for human consumption. To avoid excessive residues on fodder and forage, do not apply within 7 days of harvest.

POTATOES: COLORADO POTATO BEETLE: 10 pounds per acre. Leafhopper and Flea Beetle — 20 pounds per acre. Lygus Bugs — 40 pounds per acre. Apply when insects appear. Apply dust thoroughly to upper and lower leaf surfaces. If needed, repeat applications at 7 to 10-day intervals.

TOMATOES: TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES, RUSSET MITE: 30 to 40 lbs. per acre. Tomato Hornworm, European Corn Borer, Stink Bugs — 60 to 70 lbs. per acre. May be applied up to the day of harvest. Apply thoroughly at first signs of insect infestation. Repeat as necessary.

PEANUTS: Apply at the rate of 20-30 lbs. per acre to control Thrips, Leafhoppers, Cucumber Beetles, Bean Leaf Beetle, Velvet Bean Caterpillar, Corn Earworm, Armyworm and Leaf Spot Disease. Can be used up to day of harvest. For disease control apply when plants are 40 days old and repeat every 10 - 14 days for 4 applications. Repeat applications after rains.

WARNING

SEE REVERSE SIDE

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

See other warnings on this label
USDA Reg. No. 4185 - 160

Distributed by THE BORDEN CHEMICAL COMPANY Norfolk, Va.
SMITH-DOUGLASS Division


HARMFUL IF INHALED OR SWALLOWED on skin, or on clothing. Avoid take internally. Skin contact may be harmful. Avoid contact. ing. Take shower or bath after long sleeved work clothing. Change daily.

Note for Physician: Carbaryl is a cholinesterase inhibitor. Atropine

Keep out of any body of water,

For protection of honey bees avoid during periods when honey bees are present. When necessary to use during such periods, beekeepers well in advance to 1 mile distance until one week after a

Dispose of wastes by burying in from water supplies. Container should be buried with wastes or by smoke).



MIN*-SULFUR 5-75 DUST

ACTIVE INGREDIENTS:	5.00%
Carbaryl (1-naphthyl N-methylcarbamate)	75.00%
Sulfur	20.00%
INERT INGREDIENTS:	-----
TOTAL	100.00%

*Trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate.

ICIDE
ACT
STER-
BIECT

DIRECTIONS FOR USE

E LABEL. USE STRICTLY IN ACCORDANCE
CAUTIONS, WARNINGS AND DIRECTIONS.

PE LEAF FOLDER: 1st Brood: 20 to 25 lbs.
before first brood larvae emerge from the
nd and 3rd Broods: 20 to 40 pounds per
applied up to harvest. Grape Leafhopper
per acre. Thorough coverage is essential.
ion may be expected to control DDT-resist-
ers.

ICAN BEAN BEETLE: 10 pounds per acre.
— 20 pounds per acre. Lygus, Stink Bugs
s per acre. May be used up to harvest.
ughly at first signs of insect infestation.
ary repeat every 7 to 10 days. Do not feed
to meat or dairy animals.

LEARWORM: 30 to 40 lbs. per acre. Apply
rvals starting when silks first appear. Make
ations until silks dry or turn brown. May be
to harvest on sweet corn to be used for
umption. To avoid excessive residues on
forage, do not apply within 7 days of

COLORADO POTATO BEETLE: 10 pounds
hopper and Flea Beetle — 20 pounds per
Bugs — 40 pounds per acre. Apply when
ar. Apply dust thoroughly to upper and
urfaces. If needed, repeat applications at
ntervals.

TOMATOES: ~~TOMATO FRUITWORM, FALL ARMYWORM,~~
~~FLEA BEETLES, RUSSET MITE:~~ 30 to 40 lbs. per acre.
Tomato Hornworm, European Corn Borer, Stink Bugs —
60 to 70 lbs. per acre. May be applied up to the day of
harvest. Apply thoroughly at first signs of insect infesta-
tion. Repeat as necessary.

PEANUTS: Apply at the rate of 20-30 lbs. per acre to
control Thrips, Leafhoppers, Cucumber Beetles, Bean
Leaf Beetle, Velvet Bean Caterpillar, Corn Earworm,
Armyworm and Leaf Spot Disease. Can be used up to
day of harvest. For disease control apply when plants
are 40 days old and repeat every 10 - 14 days for 4
applications. Repeat applications after rains.

WARNING

SEE REVERSE SIDE

NOTICE OF WARRANTY: The Borden Chemical Company, Smith-Douglass Division, warrants only
that the contents of the container correspond to the analysis printed on the container,
label, or tag: THERE IS NO WARRANTY OF MERCHANTABILITY AND THERE ARE NO WARRANTIES THAT
EXTEND BEYOND SUCH ANALYSIS. Buyer assumes all risks of use, storage or handling, whether
in accordance with directions or not.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

See other warnings on this label
USDA Reg. No. 4185 - 160

Distributed by THE BORDEN CHEMICAL COMPANY Norfolk, Va.
Smith-Douglass Division

HARMFUL IF INHALED OR SWALLOWED. Do not get in eyes,
on skin, or on clothing. Avoid breathing dust. Do not
take internally. Skin contact with any insecticide
may be harmful. Avoid contact. Wash hands before eat-
ing. Take shower or bath after work. Wear regular
long sleeved work clothing. Change to clean clothes
daily.

Note for Physician: Carbaryl is a moderate, reversible,
cholinesterase inhibitor. Atropine is antidotal.

Keep out of any body of water, fish may be killed.

For protection of honey bees avoid use, if possible,
during periods when honey bees are visiting the crop.
When necessary to use during such periods, warn
beekeepers well in advance to locate hives at a safe
distance until one week after application.

Dispose of wastes by burying in non-crop lands away
from water supplies. Containers should be disposed
of by burying with wastes or by burning (Keep out of
smoke).