



Sevin Copper 3-7 Dust

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methylcarbamate)	3.0%
Copper Expressed as Metallic (Derived from Cuprous and Cupric Oxides)	7.0%
INERT INGREDIENTS:	90.0%
TOTAL	100.0%

*Trademark of Union Carbide Corporation for the active ingredient
1-naphthyl N-methylcarbamate.

DIRECTIONS FOR USE

Begin applications before diseases appear.

POTATOES: To control Colorado Potato Beetle, Leafhopper, Flea Beetle, Early and Late Blight: Dust at the rate of 30 to 40 lbs. per acre, depending on size of plants and severity of insect and disease conditions. Apply dust thoroughly to upper and lower leaf surfaces. Repeat applications at 7-10 day intervals.

TOMATOES: To control Tomato Fruitworm, Flea Beetle, and Early and Late Blight — Dust at the rate of 30 to 35 lbs. per acre. Repeat applications, if necessary, at 7-10 day intervals. Can be applied the day of harvest.

CUCUMBERS AND SUMMER SQUASH: To control Striped and Spotted Cucumber Beetle, Squash Bug, Flea Beetle, Pickworm, Melonworm, Downy Mildew, Anthracnose, Bacterial Wilt and Alternaria Leaf Spot — Dust at the rate of 30 to 35 lbs. per acre. Repeat applications, if necessary, at 7-10 day intervals. Can be applied the day of harvest.

PEPPERS AND EGGPLANT: To control Cercospora Leaf Spot on Peppers and Alternaria on Eggplant: Dust at the rate of 30 to 35 lbs. per acre. Repeat application, if necessary, at 7 to 10 day intervals. Can be applied the day of harvest.

WARNING

HARMFUL IF INHALED OR SWALLOWED. Do not get in eyes, on skin or on clothing. Do not take internally. Avoid prolonged or repeated breathing of dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Avoid storage near feed and food products.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

NOTICE OF WARRANTY: The Borden Chemical Company, Smith-Douglass Division, warrants only that the contents of the container correspond to the analysis printed on the container, label, or tag: THERE IS NO WARRANTY OF MERCHANTABILITY AND THERE ARE NO WARRANTIES THAT EXTEND BEYOND SUCH ANALYSIS. Buyer assumes all risks of use, storage or handling, whether in accordance with directions or not.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN
See other cautions on this label

Distributed by THE BORDEN CHEMICAL COMPANY
SMITH-DOUGLASS DIVISION • Norfolk, Va.