

Label Copy
NUTRO Fruit Tree Spray

Front Panel -

NUTRO Emblem

FRUIT TREE SPRAY

(Picture of a
Peach and Apple)

Use on Apples, Peaches, Pears,
Cherries and Strawberries

Back Panel -

NUTRO Emblem

FRUIT TREE SPRAY

DIRECTIONS: Apply with hose end or tank type sprayer. Mix NUTRO Fruit Tree Spray with water at the rate of five level tablespoonfuls per gallon, or 24 ozs. to 18 gallons of water. It is best to mix a little water with the NUTRO Fruit Tree Spray and make into a paste, then add the total amount of water before spraying. Application should be made in evening or early morning when there is little air movement. Spray plants lightly but thoroughly, covering all parts - particularly upper and lower portions of the leaves. Keep solution agitated by shaking or stirring. Do not spray fruits in full bloom.

A. BENTON WITH COMPANY

Back Panel - continued

Label Copy
NUTRO Fruit Tree Spray

The following table shows spray schedule for major fruit crops:

DUST OR SPRAY SCHEDULE IN LEVEL TABLESPOONS PER GALLON WATER					
STAGE OF TREE DEVELOPMENT	APPLE	PEACH	CHERRY	PEAR	STRAW-BERRIES
DORMANT	A	A	A	A	
GREEN TIP	5 tbs.				5 tbs.
PRE-BLOOM	5 tbs.	5 tbs.	5 tbs.	5 tbs.	5 tbs.
BLOOM	DO NOT SPRAY DURING BLOOM.				
PETAL FALL	5 tbs.	5 tbs.	5 tbs.	5 tbs.	
1st COVER	5 tbs.	5 tbs.	5 tbs.	5 tbs.	
2nd COVER	5 tbs.	5 tbs.	5 tbs.	5 tbs.	
3rd COVER	5 tbs.		5 tbs.		
4th COVER	5 tbs.	5 tbs.	5 tbs.		
5th COVER	5 tbs.				
6th COVER	5 tbs.	5 tbs.		5 tbs.	
7th COVER	5 tbs.			5 tbs.	
SEE BELOW	B	B	B	B	C

- A. NUTRO Dormant oil spray.
B. Stop spraying 30 days before harvest.
C. Do not use after fruit has set.

INSECTS AND DISEASES CONTROLLED:

APPLES: Insects - Aphids, Apple Maggot, Cankerworm, Codling Moth, Curculio, Flea Beetle, Japanese Beetle, Leafhopper, Leafroller, Oriental Fruit Tree Moth, Red Spider Mite, Tent Caterpillar, Thrips.

Disease - Scab, Brooks fruit spot, sooty blotch, flyspeck, bitter rot, black rot.

PEACHES: Insects - Aphids, Cankerworm, Codling Moth, Curculio, Flea Beetle, Japanese Beetle, Leafhopper, Leafroller, Oriental Fruit Moth, Red Spider Mite, Tent Caterpillar, Thrips.

Disease - Scab, brown rot, Rhizopus.

21, 1964

Back Panel - continued

Label Copy
NUTRO Fruit Tree Spray

CHERRIES: Insects - Aphids, Curculio, Leafhopper, Leafroller, Slugs,
Red Spider Mites, Tent Caterpillar, Thrips.

Disease - Brown rot, leaf spot, Botrytis rot.

PEARS: Insects - Aphids, Cankerworms, Codling Moth, Flea Beetle,
Japanese Beetle, Leafhopper, Leafroller, Oriental
Fruit Moth, Pear Psylla, Pear Slug, Red Spider
Mite, Tent Caterpillar, Thrips.

Disease - Scab.

STRAW-
BERRIES: Insects - Aphids, Flea Beetle, Japanese Beetle, Leafroller,
Red Spider Mites, Spittlebug, Strawberry Leaf Beetle,
Strawberry Weevil, Thrips.

Disease - Leaf spot, Botrytis rot.

For best results, apply according to chart, starting at times indicated
for each fruit. Repeat spray applications as necessary.

CAUTION: Avoid inhalation of dust or spray. Avoid contact with skin
and eyes. Avoid contamination of feed and foodstuffs. Wash
thoroughly after using. Store out of the reach of children.
Remove any residue at harvest by washing or peeling. Dispose
of container promptly after it is empty. Harmful if swallowed.

4/18/64
4/18/64

Label Copy
NUTRO Fruit Tree Spray

Back Panel - continued

ACTIVE INGREDIENTS:

Sulfur.....	25.00%
Methoxychlor.....	10.00%
Captan (N-trichloromethylmercapto-4-cyclohexene-1,2,dicarboximide).....	6.00%
Malathion (O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate).....	6.00%
INERT INGREDIENTS:.....	53.00%

NOTICE: Neither the manufacturer nor the seller makes any representation or warranty, expressed or implied, and shall not be held responsible in any manner for personal injury, property damage, or other type of loss resulting from handling, storage, or use of this product, whether or not used in accordance with directions.

____ NET WEIGHT

Packaged For

SMITH-DOUGLASS CO., INC.

Home Office - Norfolk, Virginia

Handwritten:
2/1/95
4/15/95