

Front Panel

Label Copy
BORDEN BLACK SPOT AND POWDERY
MILDEW CONTROL
Net: 8 oz.
May 15, 1970 (Rev. #2)

BORDEN
TM

\$1.75

Black Spot &
Powdery Mildew
Control

Controls Black Spot and Powdery Mildew on roses, flowers,
ornamentals and fruit trees.

USDA Reg. No. 4185-300

ACTIVE INGREDIENTS:

Captan (N-Trichloromethylthiotetrahydrophthalimide)...	30.00%
Dinitro (1-Methylheptyl) phenyl crotonate*	2.70%
Other Nitrophenols and derivatives chiefly Dinitro (1-Methylheptyl) phenol*30%
INERT INGREDIENTS:	67.00%
TOTAL	100.00%

*Karathane


NET WEIGHT 8 OUNCES

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS.
See back panel for additional cautions.

Side Panels

BORDEN
TM

Black Spot &
Powdery Mildew
Control



OUR PLEDGE TO YOU -- If you are not pleased with the performance of this Borden
turf and garden product, your dealer will cheerfully replace it.

Back Panel

DISEASES CONTROLLED: Roses -- Black Spot and Powdery Mildew. Flowers and Ornamentals
-- Powdery Mildew and certain leaf spots of asters, larkspur, begonia, luhlia, carnation,
chrysanthemum, calendula, delphinium, snapdragon, hydrangea, and euonymus.

Back Panel continued

Label Copy
BORDEN BLACK SPOT & POWDERY
MILDEW CONTROL
Size: 8 oz.

DIRECTIONS

Apply with hose-end or tank type sprayer. Mix with water at the rate of TWO TABLE-
SPOONS PER GALLON OF WATER. It is best to mix with a little water, making a thin
paste, then adding total amount of water before spraying. Application should be
made in evening or morning, when air is still. Spray plants lightly but thoroughly,
covering all parts, particularly upper and lower portions of leaves. Keep solution
agitated by shaking or stirring. Repeat every 7 - 10 days and after each rain.

APPLES & PEARS: Scab, Brooks fruit spot, sooty blotch, flyspeck, bitter rot and
powdery mildew. Begin applications when flower buds swell in the spring. Repeat
at 7 - 10 day intervals and after each rain. Do not apply within 21 days of harvest.

CANTALOUPE, CUCUMBER AND SQUASH: Leaf Spot, anthracnose, downy and powdery mildew.
Do not apply within 7 days of harvest. Begin applications before diseases are ex-
pected to appear. Repeat at 5 - 7 day intervals.

INSECT AND DISEASE PROTECTION FOR ROSES AND FLOWERS: Mix 2 tablespoons Borden Broad
Control Insect Spray and 2 tablespoons Borden Black Spot and Powdery Mildew Control
per gallon of water. Use weekly to control disease and insects such as aphids, mites
and thrips. Begin application when leaves first appear in the spring. Repeat weekly.


TURF: To control brown patch and powdery mildew, use one half pound (contents of
this box) in 15 gallons of water. Apply one gallon per 100 square feet. Repeat at
7 to 10 day intervals until control is obtained.

CAUTION

Avoid inhalation of dust. May cause irritation to skin. Avoid contact with skin and
eyes. Avoid contamination of feed and foodstuffs. Do not apply when temperature is
above 90 degrees during the day or when foliage remains wet. Wash thoroughly after
use. Store out of reach of children and pets. This product is poisonous to fish.
Keep out of any body of water. Harmful if swallowed.

NOTICE: Seller warrants that this product conforms to its chemical description and
is reasonably fit for the purposes stated on the label when used in accordance with
directions under normal conditions of use but Seller makes NO WARRANTY, EXPRESS OR
IMPLIED, EXCEPT AS HEREIN STATED. Seller's liability is limited to the purchase
price of the product and this shall be buyers only remedy.

SMITH-DOUGLASS/DIVISION OF BORDEN CHEMICAL, BORDEN INC
COLUMBUS, OHIO 43203 NORFOLK, VIRGINIA 23501



Borden Incorporated
Chemical Division/Smith Douglass
Post Office Box 6
Columbus, Ohio 43203

87 JAN 1977

BORDEN BLACK SPOT & POWDERY MILDEW CONTROL
EPA Reg. No. 4185-300
Your application of May 25, 1970

1. The words "ACTIVE INGREDIENT(S)" and "INERT INGREDIENT(S)" should be printed in type of the same size and be given equal prominence. The words "ACTIVE INGREDIENT(S)" and "INERT INGREDIENT(S)" should be aligned to the same margin.

2. The preferred chemical name is 2-(1-Methylheptyl) 4,6-dinitrophenyl crotonate.

3. Request your supplier(s) to submit a statement specifically authorizing this Division to use their confidential formula(s) in support of your product's registration. Refer to PR Notice 67-3. Alternatively, request your supplier to submit to this Division their complete formula(s) giving the names and percentages of both the active and the inert ingredients for Stauffer "50% CAPTAN", and Rohm & Haas "WD KARATHANE."

4. Add a container disposal statement similar to the following:

"Do not reuse container. Destroy when empty."

2

5. Separate and give precedence to the precautionary statements pertaining to human safety.

6. The general fish, wildlife and environmental cautions must appear on the label in a paragraph set apart from other cautions and directions.

Sincerely,

Douglas L. Camp
Assistant Chief
Registration Branch

Enclosure

EPA:PR HTToma:obh 1/21/71

PR 18m 9-26

BORDEN
TM

\$1.75

Black Spot &
Powdery Mildew
Control

Controls Black Spot and Powdery Mildew on roses, flowers,
ornamentals and fruit trees.

USDA Reg. No. 4185-300

ACTIVE INGREDIENTS:

Captan (N-Trichloromethylthiotetrahydrophthalimide)... 30.00%
Dinitro (1-Methylheptyl) phenyl crotonate* 2.70%
Other Nitrophenols and derivatives chiefly Dinitro
(1-Methylheptyl) phenol*30%

INERT INGREDIENTS: 67.00%

TOTAL 100.00%

*Karathane


NET WEIGHT 8 OUNCES

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS.
See back panel for additional cautions.

Side Panels

BORDEN
TM

Black Spot &
Powdery Mildew
Control



OUR PLEDGE TO YOU -- If you are not pleased with the performance of this Borden
turf and garden product, your dealer will cheerfully replace it.

Back Panel

DISEASES CONTROLLED: roses -- Black Spot and Powdery Mildew, Flowers and Ornamentals
-- Powdery Mildew and certain leaf spots of asters, larkspur, begonia, dahlias, carnations,
chrysanthemum, calendula, delphinium, snapdragon, hydrangea, and euonymus;

002-68

Label Copy
BORDEN BLACK SPOT & POWDERY
MILDEW CONTROL
Size: 8 oz.

DIRECTIONS

Apply with hose-end or tank type sprayer. Mix with water at the rate of TWO TABLESPOONS PER GALLON OF WATER. It is best to mix with a little water, making a thin paste, then adding total amount of water before spraying. Application should be made in evening or morning, when air is still. Spray plants lightly but thoroughly, covering all parts, particularly upper and lower portions of leaves. Keep solution agitated by shaking or stirring. Repeat every 7 - 10 days and after each rain.

APPLES & PEARS: Scab, Brooks fruit spot, sooty blotch, flyspeck, bitter rot and powdery mildew. Begin applications when flower buds swell in the spring. Repeat at 7 - 10 day intervals and after each rain. Do not apply within 21 days of harvest.

CANTALOUPE, CUCUMBER AND SQUASH: Leaf Spot, anthracnose, downy and powdery mildew. Do not apply within 7 days of harvest. Begin applications before diseases are expected to appear. Repeat at 5 - 7 day intervals.

INSECT AND DISEASE PROTECTION FOR ROSES AND FLOWERS: Mix 2 tablespoons Borden Broad Control Insect Spray and 2 tablespoons Borden Black Spot and Powdery Mildew Control per gallon of water. Use weekly to control disease and insects such as aphids, mites and thrips. Begin application when leaves first appear in the spring. Repeat weekly.

TURF: To control brown patch and powdery mildew, use one half pound (contents of this box) in 15 gallons of water. Apply one gallon per 100 square feet. Repeat at 7 to 10 day intervals until control is obtained.

CAUTION

Avoid inhalation of dust. May cause irritation to skin. Avoid contact with skin and eyes. Avoid contamination of feed and foodstuffs. Do not apply when temperature is above 90 degrees during the day or when foliage remains wet. Wash thoroughly after use. Store out of reach of children and pets. This product is poisonous to fish. Keep out of any body of water. Harmful if swallowed.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but Seller makes NO WARRANTY, EXPRESS OR IMPLIED, EXCEPT AS HEREIN STATED. Seller's liability is limited to the purchase price of the product and this shall be buyers only remedy.

SMITH-DOUGLASS/DIVISION OF BORDEN CHEMICAL, BORDEN INC
COLUMBUS, OHIO 43203 NORFOLK, VIRGINIA 23501

